

Appendix S1. A survey of post-ejaculatory modifications to sperm (PEMS) throughout the Kingdom Animalia

CONTENTS

I. Cnidaria – Hydrozoa	4
II. Bryozoa – Gymnolaemata	5
III. Mollusca	6
(1) Bivalvia	6
(2) Gastropoda	7
(3) Cephalopoda	9
IV. Annelida – Polychaeta	10
V. Arthropoda	11
(1) Chelicerata	11
(a) Acari	11
(b) Araneae	14
(2) Crustacea	15
(a) Ostracoda	15
(b) Copepoda	17
(3) Hexapoda	17
(a) Collembola	17
(b) Archaeognatha	19
(c) Orthoptera	19
(d) Blatteria	21

24	(e) Coleoptera	22
25	(f) Hymenoptera	23
26	(g) Lepidoptera	24
27	(h) Diptera	26
28	(i) The fungus gnat (<i>Sciara coprophila</i>)	
29	(ii) Mosquitoes	
30	(iii) The fruit fly (<i>Drosophila melanogaster</i>)	
31	VI. Urochordata	31
32	VII. Craniata	32
33	(1) Osteichthyes - Teleostei	32
34	(2) Amphibia	34
35	(a) Anura	34
36	(b) Caudata	35
37	(3) Testudines	35
38	(4) Archosauromorpha	37
39	(5) Mammalia	38
40	(a) Monotremata	38
41	(b) Marsupialia	39
42	(c) Eutheria/Placentalia	40
43	References	46

44

45

46

A SURVEY OF POST-EJACULATORY MODIFICATIONS TO SPERM (PEMS) THROUGHOUT THE KINGDOM ANIMALIA

Among the taxon-specific PEMS described below, there is tremendous variation in the extent to which systems have been investigated and in the experimental tools employed. Consequently, we have a relatively sophisticated understanding of the cellular and molecular mechanisms underlying PEMS in model systems such as eutherian mammals (i.e. mouse, rat, rabbit and human) and the fruit fly *Drosophila melanogaster*. By contrast, our understanding of PEMS for the majority of taxa is restricted to what can be inferred from ultrastructural comparisons between sperm obtained from the male reproductive tract (MRT) and the female reproductive tract (FRT). In the descriptions below, we attempt to be explicit about methods and to share authors' conclusions and interpretations of their findings. We describe more generalized, taxon-specific aspects of the reproductive biology whenever deemed necessary to understand the described PEMS.

I. CNIDARIA – HYDROZOA

During the sexual phase of their life cycle, the oocytes of hydrozoans develop along a blastostyle within specialized female reproductive polyps called gonangia. Sperm are not capable of entering a gonangium until the oocytes within have matured. To enter, sperm first pass through a funnel-like aperture, then proceed to the surface of an oocyte *via* passageways lined with epithelial cells that lead to and surround each egg (O'Rand, 1974; O'Rand & Miller, 1974). During this journey, sperm undergo two PEMS that are likely (but have not been shown) to be causally linked. First, membrane-bound 'pro-acrosomal' vesicles are progressively lost from the apical and lateral regions of the sperm head as they move through the passageways and

putatively interact with epithelial cells. By the time sperm reach the egg, nearly 90% of vesicles have been lost (O’Rand & Miller, 1974). Second, the sperm become capacitated (i.e. fertilization competent). O’Rand (1972, 1974) experimentally demonstrated the role of epithelial cells surrounding eggs in mediating PEMS. The exposure of egg packets to sperm results in efficient fertilization, but not when eggs are first stripped of their surrounding epithelial cells. Similarly, treatment of intact packets with trypsin prior to the application of sperm also inhibited fertilization, presumably due to the loss of epithelial interaction sites. Interestingly, sperm that had passed through trypsin-treated epithelial cells were incapable of fertilization, yet subsequent exposure of these same sperm to non-trypsin-treated epithelial cells fully restored fertilization capacity.

II. BRYOZOA – GYMNOAEMATA

The reproductive strategy of the gymnolaemate bryozoan *Membranipora membranacea* is unique among animals in being the only known spermcaster with conjugated sperm (Temkin, 1994; Temkin & Bortolami, 2004). These conjugates are spermatodesms, each containing 64 ‘sibling’ sperm from each cyst that remain associated at the end of spermatogenesis. The sperm of *M. membranacea* may exhibit two different PEMS. First, conjugates dissociate after entering females, after which sperm migrate individually to the surface of ovaries (Temkin & Bortolami, 2004). Second, sperm behaviour is modified after leaving the male and again after contacting a female (Temkin, 1994; Temkin & Bortolami, 2004). Specifically, conjugates exhibit three types of waveforms: small amplitude, large amplitude and reverse (i.e. propagations proceed from the tip to the base of axonemes). The conjugates are motile within the paternal coeloms, become largely quiescent after being spawned, and then recover motility after being drawn into the

lophophores of conspecifics by colony feeding currents. Moreover, the event frequencies and durations of the different waveforms differ between these locations. Experimental evidence suggests that the behavioural changes exhibited by spermatodesms after reaching females are a response to specifically contacting conspecific maternal tissue (Temkin & Bortolami, 2004).

III. MOLLUSCA

(1) Bivalvia

Whereas most species of bivalve molluscs are gonochoristic broadcast spawners with external fertilization, smaller-bodied species within many families exhibit internal fertilization and use the mantle cavity as a brood chamber (Sastry, 1979). Some of these species are self-fertilizing hermaphrodites, but others are spermcasters and have some form of prolonged sperm storage. Clams of the genus *Mysella* brood young in their suprabranchial chambers and store sperm through attachment to specialized regions of the gill lamellae and the gill suspensory membranes (Ó Foighil, 1985b). A fine-structure study of sperm transfer and storage in *M. tumida* revealed a PEMS related to gill attachment. The sperm become densely packed with their heads facing the gill filaments. Using light microscopy to examine living gills with attached sperm, Ó Foighil (1985b) observed the sperm to be continually buffeted by powerful water currents passing through the gill ostia. Stored sperm were further observed to be actively beating their flagella. Sperm nevertheless remained firmly attached to the gills. Detailed study revealed that attachment is achieved by fine, thread-like extensions of the periacrosomal plasmalemma interdigitating with shorter, stouter gill filament microvilli (see Fig. 2C, D). No fusions of the two types of microvilli were observed but there was close apposition between their respective glycocalices, with postulated glycoprotein crosslinking (Ó Foighil, 1985b). The interpretation of the sperm

generating head microvilli as an example of PEMS is supported by a fine-structure study of sperm production and release in *M. tumida* (Ó Foighil, 1985a). Curiously, sperm in testes have no microvilli on their heads, but do have microvilli similar in appearance emanating from the midpiece. It is not until after they have been packaged into spermatophores and released through the excurrent siphon that the location of microvilli changes (i.e. they disappear from the middle piece and appear on the head; Ó Foighil, 1985a,b).

(2) Gastropoda

All internally fertilizing gastropods that have had their sperm examined before and after insemination appear to share some general features associated with sperm storage and associated PEMS, irrespective of subclass (Prosobranchia, Opisthobranchia, Pulmonata). The descriptions share features in common with that of the polychaete worm *Spirorbis spirorbis* and the bivalve *M. tumida*. The seminal receptacle of prosobranch snails, derived from a differentiated portion of the renal oviduct, is specialized for prolonged sperm storage, with sperm embedding in the epithelial cells (Fretter, 1953; Webber, 1977). A fine-structure study of the gastropod *Cochlostoma montanum* revealed that the inner wall of the seminal receptacle is composed of two cell types, both secretory. Stored spermatozoa insert their heads into invaginations at the apex of both cell types, and develop long slender digitations projecting from the periacrosomal plasma membrane extending towards (but not fusing with) the epithelial cells. The gap is filled with fibrillar material (Giusti & Selmi, 1985).

Fine-structure studies of sperm storage in the pulmonate snails *Oxyloma elegans* (Selmi, Bigliardi & Giusti, 1989) and *Arianta arbustorum* (Bojat, Sander & Haase, 2001) similarly found that sperm have their heads embedded within the membranous and microvilli-covered

invaginations of the epithelial cells of the seminal receptacle. However, in contrast with *M. tumida* (Ó Foighil, 1985b) and *C. montanum* (Selmi *et al.*, 1989), neither pulmonate snail species was observed to develop microvilli/digitations on their sperm heads. Both species did, however, exhibit other PEMS. In *O. elegans*, the mature sperm found in the seminal vesicles of the male portion of the genital duct have the unique feature of a paracrystalline body encircling the apical portion of the nucleus. Sperm that have embedded in epithelial cells of a recipient's seminal receptacle, however, have heads that are slender in appearance and have lost the paracrystalline body (Selmi *et al.*, 1989). The sperm of *A. arbustorum* lose their perinuclear sheath within the seminal receptacle and, perhaps as a consequence, the conformation of the acrosome is altered. Sperm in the hermaphrodite duct prior to insemination have an acrosome that is positioned perpendicular to the longitudinal axis of the nucleus, whereas the acrosomal and nuclear axes are nearly confluent for sperm in the seminal receptacle (Bojat *et al.*, 2001). This example of PEMS may be triggered by secretions from the epithelial cells, and increased secretory activity of these cells appears to be triggered by the arriving sperm (Bojat *et al.*, 2001).

An examination of reciprocal sperm transfer and storage in the opisthobranch *Phyllaplysia taylori* similarly observed sperm with their heads embedded in the epithelial cells of the recipient's seminal receptacle (Beeman, 1972). Although limited evidence is provided, Beeman (1972, 1977) suggests that PEMS in this taxon include substantial modifications to the sperm plasma membrane while in storage.

Some gastropods also exhibit sperm conjugation (see Section IV.2 of main paper), in some cases combined with sperm heteromorphism (for definition and description, see Section V.3g of this appendix). In gastropod species with heteromorphic sperm conjugates, the tiny fertilizing eupyrene sperm attach to giant, highly modified oligopyrene sperm, which are hypothesized to

function as ‘mobile penises’ delivering fertilizing sperm to the site of sperm storage within the recipient (Fretter, 1953). After arriving at their destination, the eupyrene sperm must detach from the oligopyrene sperm and then attach to the recipient epithelium.

(3) Cephalopoda

Copulation by cephalopods includes the transfer of spermatophores by a modified tentacle of the male, the hectocotyliized third right arm, into the oviduct of the female. After the spermatophores explosively erupt, peristaltic contractions of the oviduct move sperm into the paired oviducal glands (Mann, Martin & Thiersch, 1970; Mann, 1984), where sperm can be stored for months with their heads embedded in the epithelial cells of the spermathecae located within the glands (Froesch & Marthy, 1975). In the octopus, *Octopus vulgaris*, the epithelial cells have cilia and microvilli, yet analysis of fine-structure images suggests these components do not function in sperm attachment. The epithelial cells also lack any preformed invaginations to receive sperm. Rather, it is hypothesized that each sperm ‘drills’ into its host epithelial cell using its screw-shaped acrosome (Froesch & Marthy, 1975; Tosti *et al.*, 2001). After embedding, the sperm are inactivated until the time of ovulation (Froesch & Marthy, 1975). PEMS in *O. vulgaris* include the loss of most of the acrosomal membrane that is present in sperm within spermatophores, but absent from sperm in spermathecae. Loss of the membrane exposes the screw-shaped acrosome (see Fig. 3; Tosti *et al.*, 2001). Tosti *et al.* (2001) provide experimental evidence that progesterone secreted by the female triggers the PEMS and that sperm from spermatophores possess receptors for progesterone on their plasma membranes.

IV. ANNELIDA – POLYCHAETA

PEMS have been described for two species of marine polychaete worms: *Pisione remota* (family Pisionidae; Alikunhi, 1951; Westheide, 1988) and *Spirorbis spirorbis* (family Serpulidae; Daly & Golding, 1977; Picard, 1980). Despite having extremely different reproductive systems and sperm morphologies, these species share critical aspects of sperm–FRT interactions and in the PEMS exhibited.

Pisione remota has discrete sexes and occupies coarse sand in the shallow intertidal zone. Following direct transfer to females through copulation, the aflagellate sperm are stored within the female's receptacula seminis (Alikunhi, 1951). After arriving in this storage organ, each rod-shaped sperm sheds a vacuole-like structure and the nucleus expands greatly, and the plasma membrane differentiates to grow numerous filliform extensions that envelope apical projections of the females receptacle cells, which were interpreted as relating to long-term sperm storage (Westheide, 1988).

By contrast, *Spirorbis spirorbis* is a simultaneous hermaphrodite that is sedentary and occupies spiral, calcareous tubes. They produce flagellated sperm and are believed to be 'spermcasters' (i.e. sperm that are released into the open water column enter other individuals in the feeding current, followed by internal fertilization; Daly & Golding, 1977). After entering the spermatheca, the sperm heads embed deeply into the spermathecal cells, with each head enclosed by a filiform extension of the spermathecal cell cytoplasm. PEMS involve the development inside the spermatheca of three different membrane specializations for contact with the spermathecal cells: (1) the sperm plasmalemma forms major digitate processes that radiate into the host cell cytoplasm from approximately half-way up the acrosome (see Fig. 2A), (2) minor digitate processes radiate from around the tip of the acrosome, and (3) adjacent to the sperm

nucleus, the membranes of the sperm and spermathecal cell appear thickened by adherence of electron-dense material, with the intracellular space bridged by dense filaments to form scalariform junctions (see Fig. 2A, B; Daly & Golding, 1977; Picard, 1980).

V. ARTHROPODA

(1) Chelicerata

(a) Acari

The dramatic PEMS of ticks belonging to the family Ixodidae have been the subject of investigation since 1906 (Reger, 1962 and references therein). The sperm of all ticks are aflagellate, yet elongate (up to 1000 μm in some argasid species; Rothschild, 1961) and are motile (Reger, 1974). Motility is attributable to unique cellular processes exposed on the leading tip of the mature sperm and possibly existing as long parallel ridges (Rothschild, 1961; Oliver & Brinton, 1971). Three different kinds of sperm movement have been described: writhing movements restricted to the anterior end, anterior–posterior serpentine contortions and steady gliding movement (Oliver & Brinton, 1971). Whereas details of the PEMS, sperm storage and the means by which sperm encounter oocytes can differ substantially among taxa, the general process is similar, with sperm being inseminated as ‘prospermia’ and then undergoing dramatic remodelling, elongation and the activation of motility inside the female within a few hours after insemination (Brinton, Burgdorfer & Oliver, 1974; Mothes & Seitz, 1981).

Fine-structure studies provide detailed descriptions of the process of spermiogenesis/PEMS (within both male and female) for the argasid tick, *Amblyomma dissimili* (see Fig. 4; Reger, 1961, 1962, 1963), the ixodid tick, *Dermacentor andersoni* (Brinton *et al.*, 1974) and the tetranychid tick, *Tetranychus urticae* (Mothes & Seitz, 1981). The inseminated sperm of

231 *Tetranychus urticae* are small, amoeboid cells that lack organelles and a nuclear membrane.
 232 They are additionally unusual in possessing a double surrounding membrane: the inner one is the
 233 sperm plasma membrane and the outer one is believed to be of somatic origin. Within the
 234 receptaculum seminis of the female, the sperm develop what appear to be microtubules beneath
 235 the inner membrane. Sperm then increase in size tenfold through the decondensation of
 236 chromatin and infiltration of cytoplasmic material, presumptively from the female (Mothes &
 237 Seitz, 1981).

238 The inseminated sperm of *Dermacentor andersoni* are tubular and double in length within
 239 the female (Brinton *et al.*, 1974). In all cases, it is not a stretch of the imagination to paraphrase
 240 the complex process of sperm remodelling within the female of ixodid and argasid ticks as sperm
 241 cells turning inside out (see Fig. 4; Shepherd, Levine & Hall, 1982*a*). Oliver & Brinton (1971, p.
 242 734) describe the process as follows: “Capacitation involves sub-terminal rupturing of the
 243 peripheral-most membranes at the pointed tip of the elongated spermatid, an extension of the
 244 inner core out of the newly open end of the outer sheath with simultaneous sliding of the outer
 245 sheath back over the inner core and eventually turning in and contributing to the base of inner
 246 core at the opposite end of the cell. Finally, the entire outer sheath turns in and carries the long
 247 fusiform nucleus with it. At this stage of development, the cell processes (probably functioning
 248 in locomotion) are exposed on the exterior surface of the spermatozoon and extend the length of
 249 the cell.”

250 The bizarre PEMS of ticks may relate to the unusual interaction that takes place between
 251 sperm and the FRT as part of their unique system of fertilization. Throughout the animal
 252 kingdom, there are many instances of sperm binding to or embedding in the epithelium of the
 253 FRT (Pitnick *et al.*, 2009*b*). However, the only taxa we are aware of in which sperm regularly

penetrate and reside within female epithelial cells (as opposed to embedding in the cell membrane) are in some scale insects (Robison, 1970) and ticks. In both *T. urticae* and *D. andersoni*, sperm are observed to enter epithelial cells lining both the ducts and lumen of the ovaries (Brinton *et al.*, 1974; Mothes & Seitz, 1981). Fine-structure observations of *D. andersoni* suggest that penetration of the female epithelial cells by a sperm is critical to its gaining access to the micropyle of an oocyte (Brinton *et al.*, 1974).

The extent to which the morphogenesis and activation of tick sperm is male *versus* female mediated is somewhat unclear. The enlargement of *T. urticae* sperm within the female appears dependent on provisioning of cytoplasm from the epithelial cells inhabited by the transforming sperm. By contrast, *in vitro* experiments with sperm of the ixodid tick, *Dermacentor variabilis*, and the argasid, *Ornithodoros moubata*, by Shepherd *et al.* (1982a) and Shepherd, Oliver & Hall (1982b) have shown that the transformation occurs in two stages: (1) rupture of an operculum at one end of the prospermium and (2) subsequent eversion, elongation and activation. Both stages were successfully triggered and fully executed by exposure of prospermia to polypeptides produced by the male accessory glands that are normally added to the prospermia during ejaculation (Shepherd *et al.*, 1982a,b). Interestingly, exposure of prospermia from either species to accessory gland secretions of the other species failed to trigger the transformation of prospermia (Shepherd *et al.*, 1982b).

(b) *Araneae*

At the end of spermiogenesis within the testes, the sperm of spiders coil and then are encysted within a proteinaceous sheath that is probably secreted by epithelial cells of the deferent duct of the testes. Each small, immotile sphere may contain a single sperm ('cleistospermia') or they

may be conjugated into various-sized groups ('coenospermia'; Alberti, 1990, 2000; Michalik, Haupt & Alberti, 2004; Michalik, 2007). At the extreme, individual coenospermia of *Liphistius cf. phuketensis* can include more than 30 sperm (Michalik, 2007). In addition to sperm, the lumen of the sheath contains other secretory products of the testes or vas deferentia (Michalik, 2007). Sperm arrive in the female spermathecae in the encapsulated state and remain so for hours to years depending on the species (Foelix, 2011). A fine-structure analysis of the garden spider, *Argiope bruennichi*, revealed that sperm in the spermathecae are not morphologically different from those in the deferent duct of males (Vöcking, Uhl & Michalik, 2013). The PEMS of spiders thus include decapsulation, uncoiling and activation of sperm motility, with successful PEMS shown to be critical to fertilization (Brown, 1985; Vöcking *et al.*, 2013). Vöcking *et al.* (2013) provide evidence that decapsulation and uncoiling of sperm are two morphologically and temporally discrete events, however their ultrastructural analysis could not establish whether sperm activation was synonymous with uncoiling.

By varying receipt of sperm relative to female developmental timing (i.e. mated immediately after the final moult *versus* later in adulthood), Brown (1985) experimentally demonstrated with the golden-orb-weaving spider, *Nephila clavipes*, that the timing of PEMS was influenced by the physiological state of the female (i.e. PEMS occurring in approximately 18 *versus* 7 days, respectively) and not by maturational differences among males or by the amount of time sperm were stored in the males' pedipalps. Details of sperm–female interactions underlying the PEMS of spiders remain unknown (Herberstein, Schneider & Michalik, 2011). However, the structural complexity of the glandular epithelium of the spermathecae (Uhl, 1994, 2000; Michalik *et al.*, 2005) and the identification of multiple structural types of secretions associated with sperm within the spermathecae (Vöcking *et al.*, 2013) has contributed to a model

of females contributing different secretory products at different times, thus providing a “cascade of triggers for decapsulation and uncoiling of sperm” (Herberstein *et al.*, 2011, p. 693). It should be kept in mind, however, that male secretions present in seminal fluid have also been observed to be associated with sperm within the spermathecae and may contribute to PEMS (Burger *et al.*, 2006; Vöcking *et al.*, 2013).

(2) Crustacea

(a) Ostracoda

All non-maxillopod crustacea have aflagellate sperm that are either immotile, or their motility is poorly investigated/understood (Jamieson, 1987; Morrow, 2004). Among these, ostracods of the family Cyprididae are unusual in having secondarily evolved filiform, motile sperm. Because such sperm tails derived from an aflagellate ancestor, these tails are aflagellate, with motility accomplished by undulatory waves generated by contractile bands or by unusual membranous organelles (Gupta, 1968; Reger, 1970). The length of sperm varies greatly among species with some having evolved particularly gigantic sperm (range among 51 species: 268–11,787 μm ; Smith *et al.*, 2016).

For species examined to date, the sperm of cypridoidean ostracods are immotile in the male and motile within the female seminal receptacle (Matzke-Karasz, Smith & Heb, 2017). Sperm activation in the female has been associated with shedding of a sperm coat inside the seminal receptacle (the female organ is often observed to be filled with empty sperm coats; Wingstrand 1988; Matzke-Karasz *et al.*, 2017). However, a recent and rigorous investigation of the ostracod *Mytilocypris mytiloides* reveals that the outer coat (‘fibrous coat’ of Gupta, 1968; ‘deciduous coat’ of Wingstrand, 1988) of sperm is not moulted; rather, it is composed of

granular material (likely adhered to sperm in the male vas deferens) that is slowly removed during storage in the female. The outer coat is first noticeably less compact after about 5 h in storage, with no traces visible after 15–24 h in storage (Matzke-Karasz *et al.*, 2017; also see Gupta, 1968). The timing of dissolution of the outer coat is coincident both with sperm becoming motile and with the initiation of egg fertilization (Matzke-Karasz *et al.*, 2017). Matzke-Karasz *et al.* (2017) postulate that the adaptive value of ostracod PEMS may be to (1) adhere adjacent sperm to one another for a more organized insemination and sperm-storage process, (2) suppress motility in the male/activate motility in the female, and/or (3) facilitate the transport and release of bioactive molecules from the male to the female's seminal receptacle. Finally, Matzke-Karasz *et al.* (2017) show that the empty sperm coats frequently observed within females are likely the inner coats of sperm that died and deteriorated within the female.

(b) Copepoda

Sperm of the copepod *Tisbe holothuriae* are reported to shed their cell coat (i.e. glycocalyx) within the antrum of the female (Pochon-Masson & Garagozlou-van Ginneken, 1978, cited in Ndiaye, Mattei & Thiaw, 1997). However, given the current interpretation of this phenomenon in ostracods (see above; Matzke-Karasz *et al.*, 2017), any present interpretation of the mechanism of glycocalyx loss should be considered with caution.

(3) Hexapoda

(a) Collembola

Collembola are small, soil-dwelling non-insect hexapods with indirect sperm transfer. Males deposit spermatophores in the soil, which females later pick up. By examining sperm from ten

species representing eight genera from four families, Dallai *et al.* (2004) identified two unique (and possibly functionally related) attributes of sperm contained within the spermatophores: (1) a central, extracellular cavity containing testicular secretions, and (2) a ‘peduncle’. Early in spermiogenesis, a cytoplasmic sleeve forms beneath the sperm head. As flagellar morphogenesis progresses, the sleeve expands to form a large extracellular cavity, the lumen of which progressively fills with a secretion from the epithelial wall of the testis. As the flagellum elongates, it winds to form a lenticular disc surrounding the extracellular mass, all contained within a common plasma membrane. Only those portions of the flagellum extending away from the central cavity are observed to have their own plasma membrane. Once the process of sperm winding is complete, the final steps of sperm maturation within the testes include flattening of the rolled sperm and condensation of the material stored in the central cavity (see Fig. 6).

The peduncle is a separate extracellular structure unique to collembolan sperm (Dallai, 1970). It is a long, thin, cylindrical structure that adheres to the acrosome and protrudes away from the coiled sperm cell. Length of the peduncle varies among species, but it can exceed the total length of the sperm (see Fig. 6; Fanciulli *et al.*, 2017). Similar to the central cavity formed by the coiled sperm, the peduncle is formed late in spermatogenesis by secretions from testis epithelial cells. Biochemical analyses reveal a composition rich in glycoproteins.

Sperm cells enter the female spermathecae in a rolled and immotile state with intact peduncles (Dallai *et al.*, 2004; Döring, 1986). While stored in the female, the peduncles disappear and the sperm unravel to adopt a filiform shape, thereby releasing the contents of the extracellular cavity into the spermathecae, and they become motile (Dallai *et al.*, 2004).

Although the triggers for the unique PEMS of collembola are unknown, Dallai *et al.* (2003, p. 311) reveal unique specializations of the membrane surrounding the extracellular cavity that they

interpret as “preferential sites for receiving and transmuting environmental signals, especially whatever signal(s) induce these spermatozoa to transform into filiform and motile cells upon reaching the female spermatheca”. Among several alternative hypotheses for the adaptive value of the peduncle, Fanciulli *et al.* (2017) suggest that enzymes released from the peduncle may activate the sperm unrolling process. Based on analyses of the fine structure of the spermathecae and female accessory reproductive glands in one species of collembolan, Dallai, Zizzari & Fanciulli (2008) further postulate that secretions of the spermathecal epithelial cells trigger the dissolution of the peduncle.

(b) *Archaeognatha*

For two species of the primitively wingless jumping bristletails, *Machilis distincta* and *Machilinus kleinbergi*, the sperm at insemination have been shown to have an unusual conformation with the flagellum bent like a hairpin within a common plasma membrane (Dallai, 1972). In *M. distincta*, PEMS do not occur until the immotile sperm have reached the spermatheca, at which point the common membrane is lost, essentially doubling the length of the sperm as the flagellum uncoils and becomes motile (see Fig. 7; Dallai, 1972).

(c) *Orthoptera*

As with many other insects, the sperm of acridid and catantopid grasshoppers have a thick and somewhat rigid glycocalyx covering the entire cell surface except for the apical acrosome (Longo *et al.*, 1993; Lupetti, Mercati & Dallai, 2001). It consists of three layers of glycoprotein and is produced early during spermiogenesis (Baccetti, Rosati & Bigliardi, 1971; Yasuzumi, 1979; Lupetti *et al.*, 2001). Post-testicular modifications to the glycocalyx, including elimination

of the outermost layer and some restructuring of the innermost layer begin in the males' seminal vesicles (Lupetti *et al.*, 2001). PEMS in these species include the complete dissolution of the glycocalyx within the FRT. Studies of two different species of acridid grasshoppers provide detailed descriptions of the timing and progressive cellular changes that sperm undergo within the female (Renieri & Talluri, 1974; Longo *et al.*, 1993). Within the site of insemination (the receptaculum seminis), the glycoprotein caps of bundled sperm dissolve. A couple of hours after insemination, within the female diverticula or along the spermathecal duct, the glycocalyx progressively detaches from the underlying plasma membrane and breaks down; the total process requires 15–24 h. There also may be substantial remodelling of the plasma membrane of sperm (Renieri & Talluri, 1974; Longo *et al.*, 1993). Renieri & Talluri (1974) postulate that the dramatic PEMS of acridids are triggered by enzymes secreted by the walls of the spermathecal duct. In support of this hypothesis, Giuffrida & Rosati (1993) were able to trigger glycocalyx dissolution of acridid sperm *in vitro* by incubating sperm collected from male seminal vesicles in extract from female spermathecae [albeit the sequence of events observed and timing of the glycocalyx breakdown differed from *in vivo* observations made by Longo *et al.* (1993)]. Moreover, dissolution of the glycocalyx could only be triggered by spermathecal extract from sexually mature females; sperm were unaffected by extracts from virgin females immediately before or after moulting (Giuffrida & Rosati, 1993).

The PEMS exhibited by tettigoniid orthoptera (katydids) appear different from those of grasshoppers. The sperm of katydids are transferred within spermatophores containing numerous sperm conjugates of variable size (approximately 12–20 sperm per conjugate, depending on species; Viscuso *et al.*, 1998). These conjugates have been referred to as 'spermatodesms', but technically are 'bundles' as they are the product of secondary rather than primary developmental

mechanisms (Higginson & Pitnick, 2011). The development of sperm bundles and their ultrastructural features were carefully tracked through the reproductive tracts of both sexes by Viscuso *et al.* (1998, 2002). Four different PEMS have been described. First, within the male and at the time of spermatophore construction, the sperm within bundles have all of their heads embedded in a common mucous-like cap secreted the epithelium of the intratesticular tubule wall. While in the spermatophore, the material forming the cap is disassembled, resulting in the individualization of all sperm. *In vitro* experiments suggest that this process is mediated by products of the male accessory reproductive glands (Viscuso *et al.*, 2001). Second, sperm within the female spermathecae reform into bundles that differ structurally from those observed in males. Sperm bundles within females are much larger than those observed in males, with each containing hundreds of sperm. Sperm within female bundles are also much more tightly packed, are linked together by their acrosomes, and exhibit a heightened degree of organization, for example, by the highly ordered, parallel orientation of their acrosomes. Third, relative to sperm in bundles within females, those in males exhibit conspicuously elongated heads containing extra-chromosomal material. Finally, prior to fertilization, PEMS occur in the form of sperm bundle dissociation.

(d) *Blattaria*

Sperm of the cockroach *Periplaneta americana* obtained from the female spermathecae were found to exhibit approximately twice the beat frequency of sperm obtained from three locations within the male (testes, seminal vesicles and vasa deferentia) (Hughes & Davey, 1969). Because all sperm were tested under standardized *in vitro* conditions, the difference in beat frequency cannot be attributed to proximate interactions between sperm and FRT morphology, viscosity or

pH. The addition of extracts of either male seminal fluid or female spermathecae to suspensions of sperm from male seminal vesicles had no effect on sperm beat frequency. This result prompted Hughes & Davey (1969) to examine *in vivo* the time course of changes to sperm beat frequency following insemination. Their assay suggests that sperm beat frequency changes occur after approximately 5 h of storage within the FRT. Additional support for these observations constituting PEMS in *P. americana* comes from comparison of the ultrastructure and gross morphology of sperm from seminal vesicles with those from spermathecae. Sperm from the male did not differ from those occupying the FRT for less than 5 h. By contrast, many sperm stored in the female for 10 h and all sperm stored for 48 h exhibited conspicuous modifications to the sperm heads, including compaction of periacrosomal material accompanied by a reduction in the distance between the acrosome and the plasma membrane (Hughes & Davey, 1969).

(e) *Coleoptera*

Sperm conjugation is widespread and has evolved to take a diversity of forms among species of diving beetles (Dytiscidae; Higginson *et al.*, 2012*a,b*, 2015), whirligig beetles (Gyrinidae) (Fig. 1; Breland & Simmons, 1970; Higginson *et al.*, 2015) and ground beetles (Carabidae; Sasakawa, 2007; Takami & Sota, 2007; Schubert *et al.*, 2017). PEMS in these species obviously include conjugate dissociation (see Section IV.2 of main paper). In both whirligig beetles of the genus *Dineutus* and ground beetles of the tribes Pterostichini, Carabinae and Platynini, an unusual form of primary conjugation has evolved in which all sibling sperm from a single cyst are transferred to females with their heads embedded in a central hyaline rod referred to as a ‘spermatostyle’ (Breland & Simmons, 1970; Sasakawa, 2007; Higginson & Pitnick, 2011; Schubert *et al.*, 2017). The conjugates dissociate after reaching the female’s spermatheca, resulting in mated females

having what appear to be well-organized stacks of ‘naked’ rods within their spermatheca (S. Pitnick, personal observation).

(f) *Hymenoptera*

Some of the most sophisticated research to date on the biochemical environment encountered by insect sperm during insemination (Collins *et al.*, 2006; Baer *et al.*, 2009b), while stored in the female spermathecae (Al-Lawati, Kamp & Bienefeld, 2009; Baer *et al.*, 2009a) and on associated PEMS (Poland *et al.*, 2011) has been performed using the honeybee, *Apis mellifera*. Poland *et al.* (2011) compared the proteome of sperm in ejaculates with that of sperm obtained from the spermathecae of 9–24-month-old queens. Whereas no qualitative differences were found, 15 major sperm proteins exhibited significant differences in abundance between the two samples. Three of these proteins were structural, two were of unknown function, and the remaining ten proteins had functions related to energy metabolism. Consistent with these patterns, enzymatic assays performed for some of these proteins revealed reduced activity by stored relative to ejaculated sperm (Poland *et al.*, 2011). It has not yet been possible to discriminate the extent to which the observed changes in sperm during storage in the FRT of *A. mellifera* are attributable to sperm ‘senescence’, female-mediated changes or adaptive plasticity.

Another group of social hymenoptera, the leaf-cutter ants, have also been shown to exhibit PEMS in the form of changes to sperm motility. Using an *in vivo* assay to investigate sperm–FRT interactions in *Atta colombica*, Liberti, Baer & Boomsma (2016) showed that an extract of the FRT (specifically, the bursa copulatrix and spermatheca) dramatically increases the proportion of motile sperm and sperm velocity relative to sperm exposed to saline or extracts of female haemolymph or hindgut. Interestingly, the same *in vivo* assay applied to another leaf-

cutter ant species, *Acromyrmex echinator*, revealed that changes to sperm motility are induced by the seminal fluid of rival males, but not by exposure to self seminal fluid (Liberti, Baer & Boomsma, 2018).

(g) *Lepidoptera*

With the exception of two species in the genus *Micropterix* (suborder Zeugloptera), males of all species of butterflies and moths exhibit sperm heteromorphism (also referred to as ‘sperm polymorphism’, ‘polymegaly’ and ‘dichotomous spermatogenesis’), which is the phenomenon of males producing more than one (nearly always two, but see, e.g. Au, Reunov & Wu, 1998) distinct morphological classes of sperm through tightly regulated processes. Sperm heteromorphism has arisen independently numerous times throughout the animal kingdom (and has been the subject of numerous reviews; e.g. Swallow & Wilkinson, 2002; Friedländer, Seth & Reynolds, 2005; Till-Bottraud *et al.*, 2005; Higginson & Pitnick, 2011). The different types of sperm perform discrete functions. In virtually all known cases, only one of the sperm types (often referred to as ‘eusperm’ or ‘eupyrene’ sperm) ever functions genetically in egg fertilization. The non-fertilizing sperm type (or ‘parasperm’) may lack a nucleus (‘apyrene’ sperm), contain only a partial complement of chromosomes (‘oligopyrene’ sperm), or possess the normal chromosome complement. Irrespective of their nuclear composition, parasperm are not functional in fertilization (e.g. Buckland-Nicks, 1998; Snook & Karr, 1998; Kubo-Irie *et al.*, 2003; Hayakawa, 2007). Numerous hypotheses for the adaptive value of parasperm have been proposed (Silberglied, Shepherd & Dickinson, 1984; Swallow & Wilkinson, 2002; Friedländer *et al.*, 2005; Till-Bottraud *et al.*, 2005; Holman & Snook, 2006), with some receiving empirical support (e.g. Fretter, 1953; Cook & Wedell, 1999; Sahara & Takemura, 2003; Takemura *et al.*,

2006; Holman & Snook, 2008). However, the function of parasperm within the FRT is unknown in most instances.

In Lepidoptera, apyrene sperm comprise 11–99% of the ejaculate, depending on species (Friedländer *et al.*, 2005). The two types of sperm differ structurally in other ways. First, the surface of eupyrene sperm are far more complex and ‘decorated’ than that of apyrene sperm (Friedländer *et al.*, 2005). Second, the head cyst cell and the enclosing cyst cells are lysed during release of sperm from the testes only in the case of apyrene sperm, such that all ‘sibling’ apyrene sperm are individualized and dispersed by the time they reach the seminal vesicle. By contrast, eupyrene spermatozoa remain organized as bundles throughout the male genital duct, with a matrix of ‘wavy fibres’ binding all sperm together within each bundle. Third, whereas both sperm types are highly modified after leaving the testes, including becoming ensheathed, the characteristics of the outer envelope differ between sperm types. The envelope of individual apyrene sperm, formed by secretions of epithelial cells in the upper vas deferens (Riemann & Giebultowicz, 1991, 1992), is continuous over the entire sperm surface. For eupyrene sperm, the laciniate appendages that uniquely decorate their exterior begin to disintegrate as the cysts leave the testis, with the breakdown products possibly forming the bi-layered envelope that surrounds each spermatozoan (Riemann, 1970; Riemann & Thorson, 1971). Also unique to eupyrene sperm, there is a longitudinal slit in the envelope that is plugged by a reticulate appendage (Friedländer *et al.*, 2005).

As far as is known, all lepidopterans exhibit PEMS, including the dissociation of eupyrene bundles within the spermatophore. In *Bombyx mori*, this process has been attributed to the action of initiatorin, a trypsin-like arginine C-endopeptidase secreted by cells in the prostatic region of the male’s distal ejaculatory duct, and likely also involves other organic acids, such as succinate,

present in the spermatophore (Osanai, Kasuga & Aigaki, 1989*a*; Aigaki *et al.*, 1994; Osanai & Isono, 1997). Dissociation of eupyrene bundles is temporally coupled with the activation of both eupyrene and apyrene sperm motility. Although this appears also to involve the action of cyclic-AMP (Osanai, Kasuga & Aigaki, 1989*b*), the precise mechanisms underlying this phenomenon remain unclear (Friedländer *et al.*, 2005). Other PEMS differ between eupyrene and apyrene sperm, consistent with structural differences described above. By the time sperm have reached the female's spermatheca, the eupyrene sperm have lost their reticulate appendage, leaving behind an open slit in the envelope (Riemann, 1970; Riemann & Thorson, 1971). The eupyrene sperm shed their extracellular envelopes in the spermatheca, whereas the apyrene sperm do not (Friedländer, Jeshtadi & Reynolds, 2001). Both the apyrene sperm and any eupyrene sperm that fail to exit from their envelopes eventually degenerate (Friedländer *et al.*, 2005).

(h) *Diptera*

(i) *The fungus gnat, Sciara coprophila*

The sperm of *S. coprophila* are immotile at insemination, during transfer to storage, and throughout their first few hours in the female's paired spermathecae. Then, after approximately 5 h, the sperm become organized parallel to one another, coil and begin to undulate while undergoing dramatic ultrastructural modifications that include (1) elimination from the sperm of nearly all of the electron-transparent, non-paracrystalline component of the mitochondrial derivative, (2) repositioning of the paracrystalline rod and, extraordinarily, (3) the uncoiling and later recoiling of the unusual axial filament complex into a different (see below) configuration (see Fig. 8; Makielski, 1966; Phillips, 1966*a,b*). By approximately 7 h after entering the spermathecae, the sperm have uncoiled and the transformation is complete. The sperm, now

consistently motile, occupy the periphery of the spermathecal capsules, whereas the sloughed off ribbons of mitochondrial derivative can be observed in knots in the centre of the spermathecae (Makielski, 1966). Phillips (1966a) observed that the sloughed mitochondrial material occupies most of the volume of the spermathecae. The axial filament complex of *S. coprophila* sperm is highly unusual, consisting of approximately 70 doublet tubules, each with an associated singlet tubule and organized in a spiral, rather than the $9 + 9 + 2$ pattern typical of insect sperm (Phillips, 1966a,b; Dallai, Bernini & Giusti, 1973; Dallai, 2014). The spiral structure lies adjacent to the crystalloid mitochondrial rod near the sperm head, but more caudally and for the entire length of the cell the axial filament wraps completely around the crystalloid (Phillips, 1966a,b). The enantiomorphic form of the axial filament spiral is the same in all sperm. However, the direction of coiling in all mature sperm from the female spermathecae is invariably the mirror image of sperm from the testes (see Fig. 8; Phillips, 1966a,b). Makielski (1966) notes that the speed by which the transformation occurs varies among females, requiring about an hour longer in young relative to old females. This observation suggests the transformation is influenced by the female and not strictly attributable to ‘programmed’ changes intrinsic to sperm, prompting Makielski (1966) to postulate an activational role of the spermathecal fluid.

(ii) Mosquitoes

Events occurring between insemination and fertilization in mosquitoes, including any known PEMS, were recently reviewed (Degner & Harrington, 2016). The sperm of several species have a thick glycocalyx that is developed in the males’ seminal vesicles and is present at insemination, but begins degrading within the spermathecae and is completely removed within 24 h post-mating (Clements & Potter, 1967; B  o & de Souza, 1993; Ndiaye *et al.*, 1997). On a perhaps

unrelated note, there have been claims of sperm hyperactivation within the female's bursa copulatrix (Degner & Harrington, 2016), although this is only weakly supported by circumstantial evidence. *In vitro* studies of *Culex quinquefasciatus* show that sperm can exhibit three discrete flagellar wave patterns (Thaler *et al.*, 2013), with a similar range of motility having been reported for *Aedes aegypti* (Jones & Wheeler, 1965*a,b*). These changes in motility were demonstrated to be calcium-dependent, with the full range of expression triggered *in vitro* by exposure to seminal constituents from the male accessory reproductive organs (Thaler *et al.*, 2013). Claims of hyperactivation are based on juxtaposition of the *in vitro* observations of Thaler *et al.* (2013) with the observation that sperm behaviour changes within the FRT. Specifically, rapidly spinning aggregates have been observed to form in the bursa and at the base of the spermathecal ducts (Degner & Harrington, 2016). These behaviours may be attributable, however, to the effects of viscosity of the FRT, hydrodynamic interactions among sperm, and/or interactions between sperm and FRT architecture.

(iii) *The fruit fly, Drosophila melanogaster*

The sperm of *D. melanogaster* undergo molecular changes within the female. One molecular change is the likely fusion of exosomes from the female and/or the seminal fluid with sperm (Corrigan *et al.*, 2014). It is not yet known what these exosomes transmit to sperm in *Drosophila* but in mammals, exosomes such as prostasomes and epididymosomes are thought to transport RNAs and proteins into sperm (reviewed in Aalberts, Stout & Stoorvogel, 2014). A second molecular change to sperm is that, at least in *D. melanogaster*, certain seminal proteins bind to the sperm inside the female. Foremost among these is the sex peptide (SP), a 36-amino-acid *Drosophila*-specific seminal peptide. SP causes a multitude of major and important changes to

599 the behaviour and physiology of the mated female fly, including repressing her receptivity to
 600 remating and her sleep, and increasing her egg production, feeding, gut size, and aggression
 601 (Chapman *et al.*, 2003; Lui & Kubli, 2003; Peng *et al.*, 2005; Avila *et al.*, 2010, 2011; Cognigni,
 602 Bailey & Miguel-Aliaga, 2011; Apger-McGlaughon & Wolfner, 2013; Bath *et al.*, 2017). Some
 603 of these effects are mediated through the female's receptor for SP, a G-protein coupled receptor
 604 called sex peptide receptor (SPR) (Yapici *et al.*, 2008). Interestingly, these post-mating changes
 605 have (where tested) been shown to require the presence of sperm and persist while the female is
 606 storing sperm (Manning, 1967; Kalb, DiBenedetto & Wolfner, 1993). This led Eric Kubli and
 607 colleagues to hypothesize, and then demonstrate, that SP binds to sperm (*via* its N-terminal
 608 region) and is thus retained within the female together with sperm. SP is bound along the entire
 609 length of the sperm, suggesting that sperm might serve as a protected storage site for SP (Peng *et al.*
 610 *et al.*, 2005). SP does not bind to sperm on its own, but requires a network of approximately nine
 611 other seminal proteins (proteases, lectins, cysteine-rich secreted proteins) to mediate its binding
 612 (Ravi Ram & Wolfner, 2009; LaFlamme, Ravi Ram & Wolfner, 2012; Findlay *et al.*, 2014;
 613 Singh *et al.*, 2018). SP and the network proteins are found bound to sperm soon after entry into
 614 the female, and they enter the storage organs with the sperm. Within 1–2 days the network
 615 proteins have disappeared, presumably having modified the sperm, SP, or both to allow SP to
 616 remain bound to sperm. Over time, the C-terminal (active) portion of SP is gradually cleaved
 617 from sperm (Peng *et al.*, 2005) by a trypsin-family protease of as-yet unknown origin.
 618 Interestingly, this cleavage occurs primarily on SP bound to the sperm's tail (Peng *et al.*, 2005),
 619 suggesting that there may be molecular differences in the nature of SP binding to the tail *versus*
 620 the head of sperm, but these are unknown. The released C-terminal peptide can then bind to SPR

and cause continued post-mating changes in receptivity, egg-laying, and other phenomena, in the female (Yapici *et al.*, 2008).

Drosophila melanogaster sperm may undergo additional PEMS that modulate their motility. Support for this contention comes from two reports that focus on effects of the sperm-enriched protein Amo, which is a homolog of the human transient receptor potential channel TRPP2, encoded by the gene *PKD2* (Gao, Ruden & Lu, 2003; Watnick *et al.*, 2003). Amo is localized at sperm heads and the tips of flagella, and *D. melanogaster* males with Amo mutations produce and ejaculate motile sperm that fail to reach or to be retained in the female sperm-storage organs (Gao *et al.*, 2003; Watnick *et al.*, 2003; Köttgen *et al.*, 2011). First, the waveform of wild-type *D. melanogaster* sperm was reported to differ when they are in the bursa *versus* in the seminal receptacle (Yang & Lu, 2011), with abnormal waveforms seen in sperm lacking *PKD2*. Second, in a single investigation using transgenic *D. melanogaster* with distinct fluorescent tags on their sperm heads and flagellae, Köttgen *et al.* (2011) quantified sperm beat frequency *in vitro*. They reported that although sperm released into saline from the male's seminal vesicles of wild-type and Amo-mutant males did not differ in beat frequency, wild-type sperm released from the mated female's bursa exhibited a significantly faster beat frequency than did sperm from Amo-mutant males, suggesting a change within the FRT that depended on the presence of Amo in sperm.

Other PEMS may occur in some *Drosophila* species with giant sperm. Variation in sperm length among species of *Drosophila* exceeds that in the remainder of the animal kingdom (Pitnick, Markow & Spicer, 1995a). For some species, such as *D. bifurca* with 58,290 μm long sperm (Pitnick, Spicer & Markow, 1995b; Lüpold *et al.*, 2016), the sperm are individually rolled into balls within the MRT (Joly, Luck & Dejonghe, 2008). Shortly after insemination, the sperm

balls unravel as they enter the female's approximately 8 cm long seminal receptacle (Pitnick, Markow & Spicer, 1999). We are unaware of studies of the molecular basis for this dramatic conformational change, but we think it is likely to involve PEMS.

VI. UROCHORDATA

Tunicates exhibit a diversity of asexual and sexual reproductive systems. The colonial ascidian, *Diplosoma listerianum*, is an outcrossing hermaphrodite. They are spermcasters with sperm released from the seminal vesicles of individual zooids that leave the colony through a common exhalent opening to disperse into the open sea. Sperm enter the fertilization canals of acting female zooids. Fertilization is internal, within a cloacal chamber that is isolated from sea water (Burighel & Martinucci, 1994a,b). Sperm reach the chamber containing eggs by passing down a 'fertilization canal' formed by a hollow, single-cell thick, tube-like extension of the ovary. At the base of this canal, sperm must pass through the ovary epithelium by using their corkscrew-shaped heads to spiral through the intercellular junctions binding adjacent epithelial cells (Burighel & Martinucci, 1994b). Once in this storage location, sperm may remain viable for up to one month (Bishop & Ryland, 1991).

Sperm–female interactions in *D. listerianum* have been shown to be intimate and sophisticated, with epithelium of the fertilization duct being capable of 'assessing' sperm surface proteins and blocking passage by genetically incompatible (e.g. self) sperm to the cloacal chamber. Those sperm sharing self-recognition markers with the maternal tissue are removed *via* immune-like phagocytotic processes (Burighel & Martinucci, 1994a; Bishop, 1996; Bishop, Jones & Noble, 1996). Remarkably, this system has also been shown to favour sperm of

genotypes that are under-represented in the population (a ‘rare male effect’; Pemberton, Noble & Bishop, 2003).

The fertilization canal of *D. listerianum* is also the site of unique PEMS. All sperm from the testes or those entering the fertilization duct have an unusual head structure characterized by having the nucleus flanked by both a unique elongate mitochondrion and by endoplasmic derivatives. The head also possesses a dense groove, which is an invagination of the plasmalemma bound to the nuclear envelope and which runs spirally around the entire head (see Fig. 10; Burighel, Martinucci & Magri, 1985). In striking contrast, sperm that have successfully traversed the fertilization canal have their mitochondrion and endoplasmic derivatives restricted to the posterior half of the head, with the anterior portion having become much thinner and more needle-like. The dense groove has detached from the base of the head and transformed into a helical whorl that wraps around the head seven or eight times, giving the anterior portion of the head the appearance of a corkscrew (see Fig. 10; Burighel & Martinucci, 1994a).

Some of the cellular machinery underlying the unusual PEMS of *D. listerianum* may be homologous with the ‘sperm reaction’ of solitary ascidians with external fertilization. Upon contact with the vitelline coat of an egg, the mitochondrion is rapidly translocated from the head to the tail, from which it is then lost completely (Ursprung & Schabtach, 1965; Lambert, 1982).

VII. CRANIATA

(1) Osteichthyes – Teleostei

The majority of bony fish species reproduce by spawning, during which paired females and males coordinate their simultaneous release of gametes. Fertilization happens rapidly, and sperm tend to be short-lived relative to those of externally fertilizing species with broadcast release of

gametes (Johnson & Yund, 2004; Browne et al., 2015). Fish sperm are typically activated upon contact with water (see Section IV.1 of main paper). Their behaviour is subsequently modified by contact with ovarian fluid (coelomic fluid), a semiviscous liquid derived from the ovarian secretory epithelia. This material contains proteins, sugars, hormones and inorganic ions and is released by females with eggs and retained in a boundary layer around each egg (Lahnsteiner, Weismann & Patzner, 1995; Rosengrave *et al.*, 2009; Johnson *et al.*, 2014). Ovarian fluid acts as a chemoattractant for sperm and influences sperm performance, including increases to the per cent motile sperm, swimming speed, linearity of movement, longevity, wave amplitude and swimming efficiency, with variables generally increasing with the concentration of ovarian fluid (e.g. Turner & Montgomerie, 2002; Rosengrave *et al.*, 2009; Butts *et al.*, 2017).

Internal fertilization has arisen independently at least eight times in fish (Stockley *et al.*, 1997), thus setting the stage for enhanced PEMS mediated by sperm–female interactions in those taxa. Even with internal fertilization, eggs within the female are still bathed in ovarian fluid. In the guppy, *Poecilia reticulata*, males transfer sperm to females along a groove in the gonopodium, a fin that has been modified to function as an intromittent organ. Perhaps as an adaptation to minimize sperm loss during transfer, the sperm are packaged as bundles of variable size that later undergo PEMS in the form of dissociation. Gasparini & Pilastro (2011) identified possible additional PEMS in this species. First, using artificial insemination, they showed that unrelated males have a competitive fertilization advantage over full siblings of females. Next, using *in vitro*, computer-assisted semen analysis, they provide compelling evidence that the fertilization advantage of unrelated males is attributable to ovarian fluid-mediated PEMS. Specifically, sperm velocity, a known predictor of competitive fertilization success in this

species, was significantly influenced by relatedness, with sperm swimming more slowly in the presence of ovarian fluid from a sibling female.

(2) Amphibia

(a) Anura

The majority of frog and toad species reproduce by external fertilization. While a mating pair is in amplexus, the female releases strands of eggs within a jelly matrix and the male ejaculates onto the jelly. The jelly envelopes surrounding the eggs are deposited in 5–8 (depending on species) distinct morphological layers as the eggs pass through different regions of the oviduct (Shivers & James, 1970*b*; Jégo, Joly & Boisseau, 1980; Coppin, Maes & Strecker, 2002). Investigations of the histochemical properties of the the jelly glands across the oviduct and of the corresponding jelly layers in numerous species of frogs and toads has revealed that the jelly is mainly composed of glycoproteins of a mucin type that contains up to 50% carbohydrates (e.g. Shivers & James, 1970*b*; Coppin *et al.*, 2002). Interestingly, the carbohydrate content exhibits striking species specificity (Coppin *et al.*, 2002, and references therein).

The requirement of the jelly coat for fertilization was first established by Newport (1851). A critical role for the jelly in frog PEMS is now clearly established. Experiments with the frog, *Rana pipiens*, utilizing a fully factorial design to compare fertiization rates of oocytes with or without jelly when exposed to sperm that either had or had not been previously exposed to jelly, were the first clear demonstration that the jelly capacitates sperm (Shivers & James, 1970*a*). This conclusion was reinforced by a study showing that treatment of *R. pipiens* jelly-coated eggs with an antibody to jelly prevents their fertilization whereas antibody-treated eggs are successfully fertilized by sperm that were previously capacitated by exposure to untreated jelly (Shivers &

James, 1971). By exposing sperm of the toad *Rhinella (Bufo) arenarum* to diffusible substances from the jelly coat (referred to as “egg water”), Krapf *et al.* (2007) showed that sperm attaining a capacitated state was correlated with a loss in sperm cholesterol and an increase in protein phosphorylation of tyrosine residues. Exposure to egg water was further shown to be necessary for the acrosome reaction to occur. In the absence of jelly, sperm still bind to the vitelline envelope, but acrosomal exocytosis does not occur (Krapf *et al.*, 2009). Finally, diffusible factors from the egg jelly have been shown to regulate sperm motility. Toad and frog sperm are activated by exposure to the lower osmolarity of the fertilization environment (i.e. pond water). However, such exposure only initiates *in situ* movement. The transition to progressive movement requires exposure to egg jelly or egg water (Simmons, Roberts & Dziminski, 2009; Krapf *et al.*, 2014).

(b) *Caudata*

To date, the described PEMS of salamanders and newts are restricted to the activation of sperm motility. Most members of this order reproduce by internal fertilization, with sperm transferred to females in spermatophores and then stored quiescently within spermathecae (Wake & Dickie, 1998; Sever, 2002). The eggs are coated with discrete, concentric layers of jelly sequentially deposited along the oviduct as in externally fertilizing amphibians. Fertilization occurs within the female’s cloaca, with sperm motility activated upon contact with the jelly. An investigation of fertilization in the newt *Cynops pyrrhogaster* revealed the presence of six layers of jelly, each with some unique carbohydrate components (Watanabe & Onitake, 2002). The outermost layer contains both a sperm motility-initiating substance (SMIS; likely a 34 kDa protein) and an associated acrosome reaction-inducing substance (ARIS) (Watanabe *et al.*, 2003, 2010).

(3) Testudines

In the Chinese soft-shelled turtle, *Pelodiscus sinensis*, spermatogenesis is seasonal and, following spermiation, sperm spend many months within the male epididymis and then the female oviduct (Zhang *et al.*, 2008). By comparing the fine structure of sperm obtained from these two locations, Zhang *et al.* (2015) detail substantial modifications to sperm over this protracted time period. Sperm found in the epididymis have a large cytoplasmic droplet attached along the posterior end of the head and flanking the entire midpiece. These cytoplasmic droplets change over time, with each containing, on average, 7.0 lipid droplets and 4.1 vacuoles in the late autumn and 1.4 lipid droplets and 14.5 vacuoles in the late spring. By contrast, sperm obtained from the oviduct or from the females' sperm-storage tubules (SSTs) had no associated cytoplasmic droplets (see Fig. 11). The authors interpreted the contents of the cytoplasmic droplet as an endogenous source of energy for sperm in the epididymis (Zhang *et al.*, 2015). The structure of the mitochondria in the midpiece also differed across locations and over time. Sperm from the epididymis and those stored for only a short time in the oviduct had mitochondria exhibiting an onion-like ultrastructure with 8–15 concentric laminated membranes circling a dense substrate core. At more intermediate stages of sperm storage in the oviduct, the number of membrane layers decreased, gaps appeared between the layers, and the volume of mitochondria was reduced, resulting in a thinning of the midpiece. The midpiece thinned further still later in the season as a result of the mitochondria returning to their normal structure of a double membrane with cristae and tight packing (see Fig. 11). These modifications were interpreted as sperm transitioning to mitochondrial metabolism, allowing ATP production *via* the oxidative phosphorylation (OXPHOS) pathway prior to fertilization (Zhang *et al.*, 2015). Note that a study

of the painted turtle, *Chrysemys picta*, similarly observed a large cytoplasmic droplet containing lipid droplets associated with the midpiece of epididymal sperm, with the droplet disappearing just prior to mating and being absent from oviductal sperm (Gist *et al.*, 2002).

(4) Archosauromorpha

Post-testicular sperm maturation, including the acquisition of fertilization capacity in the FRT (i.e. capacitation; see below) is a hallmark of eutherian mammal sperm. Recent investigations in the Australian saltwater crocodile, *Crocodylus porosus*, using mammalian *in vitro* capacitation conditions, supports analogous capacitation processes in the Crocodilia (Nixon *et al.*, 2016a, 2019b). This includes the induction of sustained motility, activation of cyclic AMP (cAMP) signalling pathways and protein phosphorylation mediated by protein kinase A (PKA). Use of comparative and quantitative proteomics revealed 126 proteins that were differentially phosphorylated in capacitated *versus* non-capacitated sperm. These same proteins exhibited substantial evolutionary overlap with those implicated in mammalian sperm capacitation, and included elements of metabolic, signal transduction and cellular remodelling pathways (Nixon *et al.*, 2019b).

By contrast, experiments utilizing *in vitro* fertilization with the fowl, *Gallus domesticus*, turkey, *Meleagris gallopavo*, and Japanese quail, *Coturnix coturnix japonica*, indicate spermatozoa of these species do not require a period of capacitation within the female in order to fertilize an oocyte (Howarth, 1971; Howarth & Palmer, 1972; Nixon *et al.*, 2014). Indeed, the sperm of fowl and quail show a rapid acrosome reaction *in vitro* in the presence of an oocyte's perivitelline membrane (Horrocks *et al.*, 2000; Nixon *et al.*, 2014). Nevertheless, a recent study of *G. domesticus* identified putative PEMS in the form of small vesicles (microvillus blebs,

MvBs, released from the apical tips of epithelial cells by apocrine secretion) fusing to the plasmalemma of sperm within the sperm-storage tubules of females. Several hypotheses for the function of this PEMS were suggested, including the reversible suppression of premature capacitation and stabilization of the plasmalemma (Bakst & Bauchan, 2015).

(5) Mammalia

(a) Monotremata

Sperm of the short-beaked echidna, *Tachyglossus aculeatus*, and of the platypus, *Ornithorhynchus anatinus*, have been shown to conjugate into relatively large bundles (typically about 100 spermatozoa per bundle) during passage through the epididymis (Djakiew & Jones, 1983; Johnston *et al.*, 2007; Nixon *et al.*, 2011, 2016b). These bundles obviously must dissociate within the FRT (see Section IV.2 of main paper). Whereas the adaptive value of sperm conjugation is unknown in these species, one of several hypotheses put forward by Johnston *et al.* (2007) was that conjugation is a mechanism for limiting the premature capacitation of sperm. Whether or not monotreme sperm capacitate (in addition to merely disassociating from bundles), however, is an open question. *In vitro* studies with echidna and platypus sperm suggest that the molecular process underlying any sperm capacitation must differ from that of eutherians, in that the elevation of intracellular cAMP levels fails to increase protein phosphorylation. Thus, either cAMP acts through alternative pathways or eutherian-like capacitation does not occur in monotremes (Nixon *et al.*, 2016b).

826 (b) *Marsupialia*

827 The sperm of some Australian marsupials have been shown to undergo PEMS within the FRT,
 828 but the processes and their regulation clearly differ from those involved in capacitation in
 829 eutherian mammals, and even from New World marsupials (Bedford, 1996; Mate & Rodger,
 830 1996). Due to these differences, the first successful *in vitro* fertilization (IVF) with a marsupial,
 831 in this case with the South American grey short-tailed opossum, *Monodelphis domestica*, did not
 832 occur until 1993 (Moore & Taggart, 1993), and traditional IVF has never been achieved with an
 833 Australian marsupial, although fertilization through intra-cytoplasmic sperm injection has been
 834 accomplished (Mate *et al.*, 2000; Magarey & Mate, 2003; Richings *et al.*, 2004).

835 The capacity to fertilize an oocyte appears to be achieved by *M. domestica* sperm simply by
 836 the dissociation of the paired sperm, which can be induced *in vitro*, and may not involve the
 837 same capacitation mechanisms described for eutherian mammals (see below; Moore & Taggart,
 838 1993). Although the ability to fertilize was not tested, paired sperm of the Virginia opossum,
 839 *Didelphis virginiana*, unpair under similar culture conditions *in vitro*, and their unpairing *in vivo*
 840 within the oviduct just prior to fertilization has also been documented (Rodger & Bedford, 1982).

841 The acrosome of Australian marsupials has been described as “remarkably stable” and the
 842 acrosome reaction is not inducible by reagents effective for eutherian sperm (Sistina *et al.*,
 843 1993a,b; Rodger, 1994; Bedford, 1996; Czarny, Mate & Rodger, 2008). Whereas fertilization in
 844 eutherian mammals entails fusion of the egg membrane with the central equatorial segment of
 845 the sperm head, the sperm of Australian marsupials lack an equatorial segment (Rodger, 1994).

846 Interestingly, the marsupial sperm head shifts, relative to the tail, between a linear and a ‘T’
 847 conformation, with the latter involved in sperm–egg binding in *Sminthopsis crassicaudata*

(Bedford & Breed, 1994; see Fig. 12). Sperm appear to complete spermatogenesis in the T confirmation, subsequently transition to the linear morphology during epididymal maturation, and then adopt the T conformation again in the FRT (Setiadi, Lin & Rodger, 1997; Lin & Rodger, 1999; Mate *et al.*, 2000). Induction of the T conformation has also been linked to sperm–oviduct epithelium interactions in the brushtail possum, *Trichosurus velpecula* (Sidhu *et al.*, 1999a), and the Bennett’s wallaby, *Macropus rufogriseus rufogriseus* (Boere, Diaz & Holt, 2011). Although it has been suggested that T-conformation acquisition may directly precede capacitation (Mate & Rodger, 1996), clear links to sperm functionality have yet to be established. Although evidence is limited, it has also been claimed that FRT secretory proteins bind to sperm and enhance *in vitro* survival and motility in the brushtail possum (Sidhu *et al.*, 1999b), although further investigation is required.

(c) *Eutheria/Placentalia*

All eutherian sperm are believed to exhibit PEMS in the form of capacitation, although it is becoming apparent that capacitation may exhibit species-specific characteristics and is therefore a process likely to include a diversifying suite of PEMS. The term ‘capacitation’ originally referred to the set of requisite changes sperm must undergo after prolonged residence in the FRT to acquire fertilization capacity (Austin, 1951, 1952; Chang, 1951). These observations were based on *in vivo* experiments in non-human mammals and, at the time, the underlying mechanisms were unknown. Despite the fact that human *in vitro* fertilization efforts pre-date the original use of the term capacitation (Pincus & Enzmann, 1934), human sperm capacitation was not successfully induced *in vitro* until 1969 (Edwards, Bavister & Steptoe, 1969), rapidly followed by the Nobel award winning research that achieved human *in vitro* fertilization

(Edwards, Steptoe & Purdy, 1970). Due to the difficulty of studying capacitation *in vivo*, sperm capacitation has been and continues to be studied largely *in vitro* and our understanding of the FRT environment with regard to capacitation-related PEMS has advanced relatively little (see de Jonge, 2017, for an excellent historical perspective).

Prior to providing an overview of the PEMS involved in mammalian sperm capacitation, it should be noted that the meaning of the term ‘capacitation’, even when applied strictly to mammals, has been debated. Capacitation encompasses a diverse array of PEMS that traditionally included two primary processes: (1) hyperactivation, which is the transition to high-amplitude asymmetric flagellar beating, and (2) the acrosome reaction, which is the exocytotic event exposing the acrosomal contents, including the oocyte cumulus matrix penetrating enzyme hyaluronidase and molecules participating in gamete fusion (Chang, 1984). Others, however, excluded the acrosome reaction and defined capacitation as the processes required to prepare sperm to undergo the acrosome reaction (Bedford, 1970; Florman & Babcock, 1991). The factor distinguishing these perspectives is whether the acrosome reaction is induced during the final stages of sperm migration or whether acrosome-intact sperm pass through the oocyte cumulus matrix and experience an interaction with zona pellucida proteins that initiate the acrosome reaction (reviewed by Florman & Fissore, 2015). We adhere to the traditional view regarding the acrosome reaction as the termination of capacitation and include an overview of recent acrosome reaction studies in our discussion below. Lastly, the importance of *in vitro* studies in model organisms to our understanding of capacitation cannot be overstated, although the use of these systems as surrogates for *in vivo* studies of capacitation across mammals has recently come under scrutiny (Okabe, 2014; Kaupp & Strunker, 2016). As such, our overview of capacitation is

reliant on generalizations derived from model organism studies, although we highlight species-specific aspects of capacitation PEMS where they have been studied.

The critical challenge facing sperm biologists is linking the well-established biochemical transitions and signalling events of capacitation to the *in vivo* life history of sperm in a highly selective FRT environment. Of the hundreds of millions of sperm inseminated as part of a typical mammalian ejaculate, only a small number are estimated to arrive successfully at the surface of the egg (Williams *et al.*, 1992; Suarez & Pacey, 2006). Sperm begin this journey with highly progressive motility and migrate through the cervical mucus, which is believed to biochemically alter sperm and initiate the remodelling and removal of membrane sterols (cholesterol and desmosterol) (de Jonge, 2017). Sperm are also exposed to reactive oxygen species (ROS) produced by leukocytes, which promotes the initiation of critical capacitation signalling pathways (see below) and, importantly, are believed to inhibit the progression of immature or low-quality sperm (de Jonge, 2017). Sperm migration is then actively assisted by contractions in the uterus, and upon reaching the oviduct, progressive motility ceases and sperm form a close association with the epithelium of the isthmic region (termed the ‘sperm reservoir’; Suarez, 2002; Suarez & Pacey, 2006). Residence in the sperm reservoir provides the opportunity for capacitation to be temporally linked to ovulation, when biochemical changes (such as changes in pH) promote hyperactivation and progression to the site of fertilization (Suarez, 2008). It is currently unclear whether achieving hyperactivation is sufficient for release from the sperm reservoir or whether molecular changes in sperm–epithelium binding affinity play an integral role (reviewed in Suarez, 2016). It is suspected that changes in binding affinity could be induced by the effects of oviductal epithelial secretions on sperm and that the specific molecular interactions underlying binding could be species specific (Coy *et al.*, 2012). If empirical

evidence supporting this phenomenon become available, the dynamics of sperm release from the sperm reservoir would be a PEMS-dependent process linked to novel sperm × female molecular interactions.

The ability to mimic *in vivo* capacitation under *in vitro* conditions, in conjunction with the use of genetic tools, has been fundamental to characterizing the biochemical reactions and signalling pathways governing capacitation [see Gervasi & Visconti (2016) and Aitken & Nixon (2013) for excellent overviews]. The critical molecular event in sperm capacitation is the activation of cAMP pathways, which ultimately result in increased intracellular Ca^{2+} levels necessary for the induction of hyperactivation. The activation of cAMP occurs through the synergistic activating effects of HCO_3^- (Gadella & Visconti, 2006) and reactive oxygen species (ROS) on soluble adenylate cyclase, which converts ATP into the second messenger cAMP. Enhanced cAMP levels, in turn, activate PKA and inhibit serine/threonine phosphatases. These pathways, in conjunction with ROS-induced efflux of sterols from the sperm plasma membrane, alter sperm membrane protein and lipid composition in the sperm head (Harrison & Gadella, 2005). This membrane hyperpolarization, accompanied by cAMP-mediated increases in pH, is believed to be conserved across mammals and essential to the activation of sperm ion exchangers that modulate sperm membrane potential and intracellular increases in Ca^{2+} levels. In turn, release of Ca^{2+} from stores located at the base of the sperm head induce the stereotypical transition in hyperactive flagellum wave form (Ho & Suarez, 2003). The importance of Ca^{2+} for the initiation of mammalian hyperactivation, and ultimately the acrosome reaction, has been long recognized (Yanagimachi, 1982), and the molecular basis of this relationship is being successfully elucidated using genetic approaches (reviewed by Gervasi & Visconti, 2016).

The traditional view of the acrosome reaction postulated that this exocytotic event was induced through interactions with the zona pellucida, and more specifically zona protein 3 (Zp3), after successfully traversing the oocyte cumulus matrix (Saling & Storey, 1979; Florman & Storey, 1982; Bleil & Wassarman, 1983). However, the ability to track mouse sperm fate using transgenic green fluorescent acrosome tags (Nakanishi *et al.*, 1999) has led to the unexpected observation that most fertilizing sperm undergo the acrosome reaction prior to interactions with the zona (Jin *et al.*, 2011) and that acrosome-reacted sperm have the capacity to fertilize oocytes with intact zona (Inoue *et al.*, 2011). Consistent with this biology is the observation that acrosome-intact sperm are found in the isthmus of the oviduct but sperm in the upper isthmus and ampulla are acrosome-reacted (La Spina *et al.*, 2016; Muro *et al.*, 2016). These findings support the traditionally held view that the acrosome reaction is part of capacitation. However, it is noteworthy that sperm that have penetrated the zona pellucida, but have not fused with the oolemma, remain vigorously motile and fertilization competent (Kuzan, Fleming & Seidel, 1984; Inoue *et al.*, 2011). Nonetheless, the acrosome reaction is critical in that it results in the exposure and translocation of sperm proteins participating in sperm–oocyte interactions, including Izumo 1, which is required for sperm–egg fusion (Okabe *et al.*, 1987; Miranda *et al.*, 2009; Aydin *et al.*, 2016; Ohto *et al.*, 2016).

In addition to capacitation, other PEMS have been described for eutherian mammals. For example, bovine seminal vesicle proteins (BSPs) bind to sperm and mediate the formation of the sperm reservoir through interactions with the oviductal epithelium (Hung & Suarez, 2012). Under *in vitro* capacitating conditions, some of these proteins are either lost from sperm or cleaved. It remains unclear whether these changes occur *in vivo* or whether they are induced by the FRT. Several species of eutherian mammals also produce sperm conjugates that must

dissociate within the FRT. Examples include the sperm ‘trains’ and ‘rosettes’ of several muroid rodent species (Immler *et al.*, 2007; Monclus & Fornes, 2016) and the ‘rouleaux’ of some species of guinea pig, armadillo, squirrel and loris (Higginson & Pitnick, 2011; Monclus & Fornes, 2016).

REFERENCES

- AALBERTS, M., STOUT, T. A. E. & STOORVOGEL, W. (2014). Prostatosomes: extracellular vesicles from the prostate. *Reproduction* **147**, R1–R14.
- AIGAKI, T., KASUGA, H., NAGAOKA, S. & OSANAI, M. (1994). Purification and partial amino acid sequence of initiatorin, a prostatic endopeptidase of the silkworm, *Bombyx mori*. *Insect Biochemistry and Molecular Biology* **24**, 969–975.
- AITKEN, R. J. & NIXON, B. (2013). Sperm capacitation: A distant landscape glimpsed but unexplored. *Molecular Human Reproduction* **19**, 785–793.
- ALBERTI, G. (1990). Comparative spermatology of Araneae. *Acta Zoologica Fennica* **190**, 17–34.
- ALBERTI, G. (2000). Chelicerata. In *Reproductive biology of invertebrates, vol. 9, Part B. Progress in male gamete ultrastructure* (ed. B. G. M. Jamieson), pp. 311–388. John Wiley and Sons, Ltd., New York.
- ALIKUNHI, K. H. (1951). On the reproductive organs of *Pisone remota* (Southern), together with a review of the family Pisionidae (Polychaeta). *Proceedings of the Indian Academy of Sciences, Section B* **33**, 14–31.
- AL-LAWATI, H., KAMP, G. & BIENEFELD, K. (2009). Characteristics of the spermathecal contents of old and young honeybee queens. *Journal of Insect Physiology* **55**, 116–121.

- 984 APGER-MCGLAUGHON, J. & WOLFNER, M.F. (2013). Post-mating change in excretion by mated
 985 *Drosophila melanogaster* females is a long-term response that depends on sex peptide and
 986 sperm. *Journal of Insect Physiology* **59**, 1024–1030.
- 987 AU, D. W.-T., REUNOV, A. A. & WU, R. S-S. (1998). Four lines of spermatid development and
 988 dimorphic spermatozoa in the sea urchin *Anthocidaris crassispina* (Echinodermata,
 989 Echinoida). *Zoomorphology* **118**, 159–168.
- 990 AUSTIN, C. R. (1951). Observations on the penetration of the sperm into the mammalian egg.
 991 *Australian Journal of Scientific Research. Series B: Biological Sciences* **4**, 581–596.
- 992 AUSTIN, C. R. (1952). The ‘capacitation’ of the mammalian sperm. *Nature* **170**, 326.
- 993 AVILA, F. W, RAVI RAM, K., BLOCH QAZI, M. C. & WOLFNER, M. F. (2010). Sex peptide is
 994 required for the efficient release of stored sperm in mated *Drosophila* females. *Genetics* **186**,
 995 595–600.
- 996 AVILA, F. W., SIROT, L. K., LAFLAMME, B. A., RUBINSTEIN, C. D. & WOLFNER, M. F. (2011).
 997 Insect seminal fluid proteins: identification and function. *Annual Review of Entomology* **56**,
 998 21–40.
- 999 AYDIN, H., SULTANA, A., LI, S., THAVALINGAM, A. & LEE, J. E. (2016). Molecular architecture of
 1000 the human sperm IZUMO1 and egg JUNO fertilization complex. *Nature* **534**, 562–565.
- 1001 BACCETTI, B., ROSATI, F. & BIGLIARDI, E. (1971). The spermatozoon of Arthropoda. XIII. The
 1002 cell surface. *Journal of Ultrastructure Research* **35**, 582–605.
- 1003 BAER, B., EUBEL, H., TAYLOR, N. L., O’TOOLE, N. & MILLAR A. H. (2009a). Insights into female
 1004 sperm storage from the spermathecal fluid proteome of the honeybee *Apis mellifera*. *Genome*
 1005 *Biology* **10**, R67.
- 1006 BAER, B., HEASLEWOOD, J. L, TAYLOR, N. L., EUBEL, H. & MILLAR, A. H. (2009b). The seminal

- 1007 fluid proteome of the honeybee *Apis mellifera*. *Proteomics* **9**, 2085–2097.
- 1008 BAKST, M. R. & BAUCHAN, G. (2015). Apical blebs on sperm storage tubule epithelial cell
1009 microvilli: Their release and interaction with resident sperm in the turkey hen oviduct.
1010 *Theriogenology* **83**, 1438–1444.
- 1011 BÀO, S. N. & DE SOUZA, W. (1993). Ultrastructural and cytochemical studies of the spermatid
1012 and spermatozoon of *Culex quinquefasciatus* (Culicidae). *Journal of Submicroscopic*
1013 *Cytology and Pathology* **25**, 213–222.
- 1014 BATH, E., BOWDEN, S., PETERS, C., REDDY, A., TOBIAS, J. A., EASTON-CALABRIA, E., SEDDON,
1015 N., GOODWIN, S. F. & WIGBY, S. (2017). Sperm and sex peptide stimulate aggression in
1016 female *Drosophila*. *Nature Ecology and Evolution* **1**, 0154.
- 1017 BEDFORD, J. M. (1970). Sperm capacitation and fertilization in mammals. *Biology of*
1018 *Reproduction* **2 (suppl 1)**, 128–158.
- 1019 BEDFORD, J. M. (1996). What marsupial gametes disclose about gamete function in eutherian
1020 mammals. *Reproduction Fertility and Development* **8**, 569–580.
- 1021 BEDFORD, J. M. & BREED, W. G. (1994). Regulated storage and subsequent transformation of
1022 spermatozoa in the fallopian tubes of an Australian marsupial, *Sminthopsis crassicaudata*.
1023 *Biology of Reproduction* **50**, 845–854.
- 1024 BEEMAN, R. D. (1972). Sperm biology in anaspidean mollusks. *Echo* **5**, 19–21.
- 1025 BEEMAN, R. D. (1977). Gastropoda: Opisthobranchia. Pp. 115–179 in *Reproduction of Marine*
1026 *Invertebrates*, Vol. IV. AC Giese and JS Pearse, eds. New York: Academic Press.
- 1027 BISHOP, J. D. D. (1996). Female control of paternity in the internally fertilizing compound
1028 ascidian *Diplosoma listerianum*. I. Autoradiographic investigation of sperm movements in
1029 the female reproductive tract. *Proceedings of the Royal Society, Series B* **263**, 369–376.

- 1030 BISHOP, J. D. D, JONES, C. S. & NOBLE, L. R. (1996). Female control of paternity in the internally
 1031 fertilizing compound ascidian *Diplosoma listerianum*. II. Investigation of male mating
 1032 success using RAPD markers. *Proceedings of the Royal Society, Series B* **263**, 401–407.
- 1033 BISHOP, J. D. D. & RYLAND, J. S. (1991). Storage of exogenous sperm by the compound ascidian
 1034 *Diplosoma listerianum*. *Marine Biology* **108**, 111–118.
- 1035 BLEIL, J. D. & WASSARMAN, P. M. (1983). Sperm-egg interactions in the mouse: sequence of
 1036 events and induction of the acrosome reaction by a zona pellucida glycoprotein.
 1037 *Developmental Biology* **95**, 317–324.
- 1038 BOERE, J., DIAZ, D. E. & HOLT, W. V. (2011). Sperm motility activation, sperm heterogeneity
 1039 and sperm-female tract interactions in Bennett’s wallaby (*Macfops*
 1040 *rufogriseus*). *Reproduction Fertility and Development* **23**, 603–617.
- 1041 BOJAT, N. C., SAUDER, U. & HAASE, M. (2001). The spermathecal epithelium, sperm and their
 1042 interactions in the hermaphroditic land snail *Arianta arbustorum* (Pulmonata,
 1043 Stylommatophora). *Zoomorphology* **120**, 149–157.
- 1044 BRELAND, O. P. & SIMMONS, E. (1970). Preliminary studies of the spermatozoa and the male
 1045 reproductive system of some whirligig beetles (Coleoptera: Gyrinidae). *Entomological News*
 1046 **81**, 101–110.
- 1047 BRINTON, L. P, BURGDORFER, W. & OLIVER, J. H. JR. (1974). Histology and fine structure of
 1048 spermatozoa and egg passage in the female tract of *Dermacentor andersoni* Stiles (Acari:
 1049 Ixodidae). *Tissue & Cell* **6**, 109–125.
- 1050 BROWN, S. G. (1985). Mating behavior of the golden-orb-weaving spider, *Nephila clavipes*: II.
 1051 Sperm capacitation, sperm competition, and fecundity. *Journal of Comparative Psychology*
 1052 **99**, 167–175.

- 1053 BROWNE, R. K., KAUROVA, S. A., UTESHEV, V. K., SHISHOVA, N. V., MCGINNITY, D., FIGIEL, C.
 1054 R., MANSOUR, N., AGNEW, D., WU, M., GAKHOVA, E. N., DZYUBA, B. & COSSON, J. (2015).
 1055 Sperm motility of externally fertilizing fish and amphibians. *Theriogenology* **83**, 1–13.
- 1056 BUCKLAND-NICKS, J. (1998). Prosobranch parasperm: sterile cells germ cells that promote
 1057 paternity? *Micron* **29**, 267–280.
- 1058 BURGER, M., MICHALIK, P., GRABER, W., JACOB, A., NENTWIG, W. & KROPF, C. (2006). Complex
 1059 genital system of a Haplogyne spider (Arachnida, Araneae, Tetrablemmidae) indicates
 1060 internal fertilization and full female control over transferred sperm. *Journal of Morphology*
 1061 **267**, 166–186.
- 1062 BURIGHEL, P., MARTINUCCI GB. (1994a). Sexual reproduction in the compound ascidian
 1063 *Diplosoma listerianum* (Tunicata). I. Metamorphosis, storage and phagocytosis of sperm in
 1064 female duct. *Marine Biology* **118**, 489–498.
- 1065 BURIGHEL P. & MARTINUCCI, G. B. (1994b). Sexual reproduction in the compound ascidian
 1066 *Diplosoma listerianum* (Tunicata). II. Sperm penetration through ovary wall and evidence of
 1067 internal fertilization. *Marine Biology* **118**, 499–510.
- 1068 BURIGHEL, P., MARTINUCCI, G. B. & MAGRI, F. (1985). Unusual structures in the spermatozoa of
 1069 the ascidians *Lissoclinum perforatum* and *Diplosoma listerianum* (Didemnidae). *Cell and*
 1070 *Tissue Research* **241**, 513–521.
- 1071 BUTTS, I. A. E., PROKOPCHUK, G., KASPAR, V., COSSON, J. & PITCHER, T. E. (2017). Ovarian fluid
 1072 impacts flagellar beating and biomechanical metrics of sperm between alternative
 1073 reproductive tactics. *Journal of Experimental Biology* **220**, 2210–2217.
- 1074 CHANG, M. C. (1951). Fertilizing capacity of spermatozoa deposited into the fallopian tubes.
 1075 *Nature* **168**, 697–698.

- 1076 CHANG, M. C. (1984). The meaning of sperm capacitation. A historical perspective. *Journal of*
 1077 *Andrology* **5**, 4550.
- 1078 CHAPMAN, T., BANGHAM, J., VINTI, G., SEIFRIED, B., LUNG, O., WOLFNER, M.F., SMITH, H.K.,
 1079 PARTRIDGE, L. (2003). The sex peptide of *Drosophila melanogaster*: female post- mating
 1080 responses analyzed by using RNA interference. *Proceedings of the National Academy of*
 1081 *Sciences, USA* **100**, 9923–9928.
- 1082 CLEMENTS, A. N. & POTTER, S. A. (1967). The fine structure of the spermathecae and their ducts
 1083 in the mosquito *Aedes aegypti*. *Journal of Insect Physiology* **13**, 1825-1836.
- 1084 COGNIGNI, P., BAILEY, A.P., MIGUEL-ALIAGA, I. (2011). Enteric neurons and systemic signals
 1085 couple nutritional and reproductive status with intestinal homeostasis. *Cell Metabolism* **13**,
 1086 92–104.
- 1087 COLLINS, A. M., CAPERNA, T. J., WILLIAMS, V., GARETT, W. M. & EVANS, J. D. (2006).
 1088 Proteomic analyses of male contributions to honeybee sperm storage and mating. *Insect*
 1089 *Molecular Biology* **15**, 541–549.
- 1090 COOK, P. A. & WEDELL, N. (1999). Non-fertile sperm delay female remating. *Nature* **397**, 486.
- 1091 COPPIN, A., MAES, E. & STRECKER, G. (2002). Species-specificity of amphibia carbohydrate
 1092 chains: the *Bufo viridis* case study. *Carbohydrate Research* **337**, 121–132.
- 1093 CORRIGAN, L., REHAI, S., LEIBLICH, A., FAN, S. J., PERERA, S. M., PATEL, R., GANDY, C.,
 1094 WAINWRIGHT, S. M., MORRIS, J. F., HAMDY, F., GOBERDHAN, D. C. & WILSON, C. (2014).
 1095 BMP-regulated exosomes from *Drosophila* male reproductive glands reprogram female
 1096 behavior. *Journal of Cell Biology* **206**, 671–688.
- 1097 COY, P., GARCÍA-VÁZQUEZ, F. A., VISCONTI, P. E. & AVILÉS, M. (2012). Roles of the oviduct in
 1098 mammalian fertilization. *Reproduction* **144**, 649–660.

- 1099 CZARNY, N. A., MATE, K. E. & RODGER, J. C. (2008). Acrosome stability in the spermatozoa of
1100 dasyurid marsupials. *Reproduction Fertility and Development* **20**, 295–302.
- 1101 DALLAI, R. (1970). The spermatozoon of Arthropoda XI. Further observations on Collembola.
1102 In: Baccetti, B. (Ed.), *Comparative Spermatology*. Academic Press, New York-London, pp.
1103 276–279.
- 1104 DALLAI, R. (1972). The arthropod spermatozoon 17. *Machilis distincta* Janetsch (Insecta,
1105 Thysanura). *Monitore Zoologico Italiano* **6**, 37–61.
- 1106 DALLAI, R. (2014). Overview on spermatogenesis and sperm structure of Hexapoda. *Arthropod*
1107 *Structure & Development* **43**, 257–290.
- 1108 DALLAI, R., BERNINI, F. & GIUSTI, F. (1973). Interdoublet connections in the sperm flagellar
1109 complex of *Sciara*. *Journal of Submicroscopic Cytology* **5**, 137–145.
- 1110 DALLAI, R., FANCIULLI, P. P., FRATI, F., PACCAGNINI, E. & LUPETTI, P. (2003). Membrane
1111 specializations in the spermatozoa of collembolan insects. *Journal of Structural Biology* **142**,
1112 311–318.
- 1113 DALLAI, R., FANCIULLI, P. P., FRATI, F., PACCAGNINI, E. & LUPETTI, P. (2004). Sperm winding in
1114 Collembola. *Pedobiologia* **48**, 493–501.
- 1115 DALLAI, R., ZIZZARI, Z. V. & FANCIULLI, P. P. (2008). Fine structure of the spermatheca and of
1116 the accessory glands in *Orchesella villosa* (Collembola, Hexapoda). *Journal of Morphology*
1117 **269**, 464–478.
- 1118 DALY, J. M. & GOLDING, D. W. (1977). A description of the spermatheca of *Spirorbis spirorbis*
1119 (L.) (Polychaeta: Serpulidae) and evidence for a novel mode of sperm transmission. *Journal*
1120 *of the Marine Biology Association, UK* **57**, 219–227.

- 1121 DEGNER, E. C. & HARRINGTON, L. C. (2016). A mosquito sperm's journey from male ejaculate to
 1122 egg: mechanisms, molecules, and methods for exploration. *Molecular Reproduction and*
 1123 *Development* **83**, 897–911.
- 1124 DE JONGE, C. (2017). Biological basis for human capacitation - revised. *Human Reproduction*
 1125 *Update* **23**, 289–299.
- 1126 DJAKIEW, D. & JONES, R. C. (1983). Sperm maturation, fluid transport, and secretion and
 1127 absorption of protein in the epididymis of the echidna, *Tachyglossus aculeatus*. *Journal of*
 1128 *Reproduction and Fertility* **68**, 445–456.
- 1129 DÖRING, D. (1986). On the male reproduction biology of *Orchesella cincta* (Collembola,
 1130 Entomobryidae). In: Dallai R (ed.), *Second International Seminar on Apterygota*. University
 1131 of Siena, Siena, pp. 171–176.
- 1132 EDWARDS, R. G., BAVISTER, B. D. & STEPTOE, P. C. (1969). Early stages of fertilization in vitro
 1133 of human oocytes matured *in vitro*. *Nature* **221**, 632–635.
- 1134 EDWARDS, R. G., STEPTOE, P. C. & PURDY, J. M. (1970). Fertilization and cleavage in vitro of
 1135 preovulator human oocytes. *Nature* **227**, 1307–1309.
- 1136 FANCIULLI, P. P., MENCARELLI, C., MERCATI, D., DALLAI, R. & LUPETTI, P. (2017). The peculiar
 1137 extra-acrosomal structure of the Collembola (Hexapoda) spermatozoa. *Micron* **101**, 114–122.
- 1138 FINDLAY, G. D., SITNIK, J. L., WANG, W., AQUADRO, C. A., CLARK, N. L. & WOLFNER, M. F.
 1139 (2014). Evolutionary rate covariation identifies new members of a protein network required
 1140 for *Drosophila melanogaster* female post-mating responses. *PLoS Genetics* **10**, e1004108.
- 1141 FLORMAN, H. M. & BABCOCK, D. F. (1991). Progress toward understanding the molecular basis
 1142 of capacitation. In: Wassarman PM, editor. *Elements of mammalian fertilization: Basic*
 1143 *concepts*. Boca Ratan, FL: CRC Press. pp. 105–132.

- 1144 FLORMAN, H. M. & FISSORE, R. A. (2015). Fertilization in mammals. In: Plant TM, Zeleznik AJ
 1145 (eds). *Knobil and Neill's Physiology of Reproduction, 4th edn*. Amsterdam: Elsevier,
 1146 pp. 149–196.
- 1147 FLORMAN, H. M. & STOREY, B. T. (1982). Mouse gamete interactions: the zona pellucida is the
 1148 site of the acrosome reaction leading to fertilization *in vitro*. *Developmental Biology* **91**, 121–
 1149 130.
- 1150 FOELIX, R. F. (2011). *Biology of Spiders*. Oxford: Oxford University Press.
- 1151 FRETTER, V. (1953). The transference of sperm from male to female prosobranch, with reference,
 1152 also, to pyramidellids. *Proceedings of the Linnean Society of London* **164**, 217–224.
- 1153 FRIEDLÄNDER, M., JESHTADI, A. & REYNOLDS, S. E. (2001). The structural mechanism of
 1154 trypsin-induced intrinsic motility in *Manduca sexta* spermatozoa *in vitro*. *Journal of Insect*
 1155 *Physiology* **47**, 245–255.
- 1156 FRIEDLÄNDER, M., SETH, R. K & REYNOLDS, S. E. (2005). Eupyrene and apyrene sperm:
 1157 dichotomous spermatogenesis in Lepidoptera. *Advances in Insect Physiology* **32**, 206–308.
- 1158 FROESCH, D. & MARTHY, H-J. (1975). The structure and function of the oviducal gland in
 1159 octopods (Cephalopoda). *Proceedings of the Royal Society, Series B* **188**, 95–101.
- 1160 GADELLA, B. M. & VISCONTI, P.E. (2006). Regulation of capacitation. In C. De Jonge, & C.
 1161 Barrett, (Eds.), *The sperm cell: Production, maturation, fertilization, regeneration* (pp. 134–
 1162 169). Cambridge: Cambridge University Press.
- 1163 GAO, Z., RUDEN, D. M. & LU, X. (2003). PKD2 cation channel is required for directional sperm
 1164 movement and male fertility. *Current Biology* **13**, 2175–2178.

- 1165 GASPARINI, C. & PILASTRO, A. (2011). Cryptic female preference for genetically unrelated males
 1166 is mediated by ovarian fluid in the guppy. *Proceedings of the Royal Society, Series B* **278**,
 1167 2495–2501.
- 1168 GERVASI, M. G. & VISCONTI, P. E. (2016). Chang's meaning of capacitation: a molecular
 1169 perspective. *Molecular Reproduction and Development* **83**, 860–874.
- 1170 GIST, D. H., DAWES, S. M., TURNER, T. W., SHELDON, S. & CONGDON, J. (2002). Sperm storage
 1171 in turtles: a male perspective. *Journal of Experimental Zoology* **292**, 180–186.
- 1172 GIUFFRIDA, A. & ROSATI F. (1993). Changes in sperm tail of *Eyprepacnemis plorans* (Insecta,
 1173 Orthoptera) as a result of *in vitro* incubation in spermathecal extract. *Invertebrate*
 1174 *Reproduction and Development* **24**, 47–52.
- 1175 GIUSTI, F. & SELMI, M. G. (1985). The seminal receptacle and sperm storage in *Cochlostoma*
 1176 *montanum* (Issel) (Gastropoda: Prosobranchia). *Journal of Morphology* **184**, 121–133.
- 1177 GUPTA, B. L. (1968). Aspects of the motility in the non-flagellate spermatozoa of freshwater
 1178 Ostracods. In: *Aspects of Cell Motility, XXII Symposium of the Society for Experimental*
 1179 *Biology* (Ed. by P. L. Miller), pp. 117–129. Oxford: Cambridge University Press.
- 1180 HARRISON, R. A. & GADELLA, B. M. (2005). Bicarbonate-induced membrane processing in sperm
 1181 capacitation. *Theriogenology* **63**, 342–351.
- 1182 HAYAKAWA, Y. (2007). Parasperm: morphological and functional studies on nonfertile sperm.
 1183 *Ichthyological Research* **54**, 111–130.
- 1184 HERBERSTEIN, M. E, SCHNEIDER, J. M. & MICHALIK, P. (2011). Sperm dynamics in spiders.
 1185 *Behavioral Ecology* **22**, 692–695.

- 1186 HIGGINSON, D. M., BADYAEV, A. V., SEGRAVES, K. A. & PITNICK, S. (2015). Causes of
 1187 discordance between allometries at and above species level: an example with aquatic beetles.
 1188 *The American Naturalist* **186**, 176–186.
- 1189 HIGGINSON, D. M., MILLER, K. B., SEGRAVES, K. A. & PITNICK, S. (2012a). Convergence,
 1190 recurrence and diversification of complex sperm traits. *Evolution* **66–5**, 1650–1661.
- 1191 HIGGINSON, D. M., MILLER K. B., SEGRAVES, K. A. & PITNICK, S. (2012b). Female reproductive
 1192 tract form drives the evolution of complex sperm morphology. *Proceedings of the National*
 1193 *Academy of Sciences, USA* **109**, 4538–4543.
- 1194 HIGGINSON, D. M. & PITNICK, S. (2011). Intra-ejaculate sperm interactions: do sperm cooperate?
 1195 *Biological Reviews* **87**, 249–270.
- 1196 HO, H. C. & SUAREZ, S. S. (2003). Characterization of the intracellular calcium store at the base
 1197 of the sperm flagellum that regulates hyperactivated motility. *Biology of Reproduction* **68**,
 1198 1590–1596.
- 1199 HOLMAN, L. & SNOOK, R. R. (2006). Spermicide, cryptic female choice and the evolution of
 1200 sperm form and function. *Journal of Evolutionary Biology* **19**, 1660–1670.
- 1201 HOLMAN, L. & SNOOK, R. R. (2008). A sterile sperm caste protects brother fertile sperm from
 1202 female-mediated death in *Drosophila pseudoobscura*. *Current Biology* **18**, 292–296.
- 1203 HORROCKS, A. J., STEWART, S., JACKSON, L. & WISHART, G. J. (2000). Induction of acrosomal
 1204 exocytosis in chicken spermatozoa by inner perivitelline-derived N-linked glycans.
 1205 *Biochemical and Biophysical Research Communications* **278**, 84–89.
- 1206 HOWARTH, JR. B. (1971). An examination for sperm capacitation in the fowl. *Biology of*
 1207 *Reproduction* **3**, 338–341.

- 1208 HOWARTH, JR. B. & PALMER, M. B. (1972). An examination of the need for sperm capacitation in
1209 the turkey, *Meleagris gallopavo*. *Journal of Reproduction and Fertility* **28**, 443–445.
- 1210 HUGHES, M. & DAVEY, K. G. (1969). The activity of spermatozoa of *Periplaneta*. *Journal of*
1211 *Insect Physiology* **15**, 1607–1616.
- 1212 HUNG, P. & SUAREZ, S. S. (2012). Alterations to the bull sperm surface proteins that bind sperm
1213 to oviductal epithelium. *Biology of Reproduction* **87**, 1–11.
- 1214 IMMLER, S., MOORE, H. D. M., BREED, W. G. & BIRKHEAD, T. R. (2007). By hook or by crook?
1215 Morphometry, competition and cooperation in rodent sperm. *PloS One* **2**, e170.
- 1216 INOUE, N., SATOUH, Y., IKAWA, M., OKABE, M. & YANAGIMACHI, R. (2011). Acrosome-reacted
1217 mouse spermatozoa recovered from the perivitelline space can fertilize other eggs.
1218 *Proceedings of the National Academy of Sciences, USA* **108**, 20008–20011.
- 1219 JAMIESON, B. G. M. (1987). *The Ultrastructure and Phylogeny of Insect Spermatozoa*.
1220 Cambridge: Cambridge University Press.
- 1221 JÉGO, P., JOLY, J. & BOISSEAU, C. (1980). Les gangues ovulaires des Amphibiens (protéines
1222 sécrétées par l’oviducte) et leurs roles dans la fécondation. *Reproduction Nutrition*
1223 *Development* **20**, 57–567.
- 1224 JIN, M., FUJIWARA, E., KAKIUCHI, Y., OKABE, M., SATOUH, Y., BABA, S. A., CHIBA, K. &
1225 HIROHASHI, N. (2011). Most fertilizing mouse spermatozoa begin their acrosome reaction
1226 before contact with the zona pellucida during in vitro fertilization. *Proceedings of the*
1227 *National Academy of Sciences, USA* **108**, 4892–4896.
- 1228 JOHNSON, S. L., VILLARROEL, M., ROSENGRAVE, P., CARNE, A., KLEFFMANN, T., LOKMAN, P. M.
1229 & GEMMELL, N. J. (2014). Proteomic analysis of Chinook salmon (*Onchorhynchus*
1230 *tshaytscha*) ovarian fluid. *PLoS One* **9(8)**, e104155.

- 1231 JOHNSON, S. L. & YUND, P. O. (2004). Remarkable longevity of dilute sperm in a free-spawning
1232 colonial ascidian. *The Biological Bulletin* **206**, 144–151.
- 1233 JOHNSTON, S. D., SMITH, B., PYNE, M., STENZEL, D. & HOLT, W. V. (2007). One-sided
1234 ejaculation of echidna sperm bundles. *The American Naturalist* **170**, E162–E164.
- 1235 JOLY, D., LUCK, N. & DEJONGHE, B. (2008). Adaptation to long sperm in *Drosophila*: correlated
1236 development of the sperm roller and sperm packaging. *Journal of Experimental Zoology:*
1237 *Series B* **310**, 167–178.
- 1238 JONES, J. C. & WHEELER, R. E. (1965a). Studies on spermathecal filling in *Aedes aegypti*
1239 (Linnaeus) I. description. *Biological Bulletin* **129**, 134–150.
- 1240 JONES, J. C. & WHEELER, R. E. (1965b). Studies on spermathecal filling in *Aedes aegypti*
1241 (Linnaeus). 2. experimental. *Biological Bulletin* **129**, 532–545.
- 1242 KALB, J. M., DiBENEDETTO, A. J. & WOLFNER, M. F. (1993). Probing the function of *Drosophila*
1243 *melanogaster* accessory glands by directed cell ablation. *Proceedings of the National*
1244 *Academy of Sciences, USA* **90**, 8093–8097.
- 1245 KAUPP, U. B. & STRUNKER, T. (2016). Signaling in sperm: more different than similar. *Trends in*
1246 *Cell Biology* **27**, 101–109.
- 1247 KÖTTGEN, M., HOFHERR, A., LI, W., CHU, K., COOK, S., MONTELL, C. & WATNICK, T. (2011).
1248 *Drosophila* sperm swim backwards in the female reproductive tract and are activated via
1249 TRPP2 ion channels. *PLoS ONE* **6**, e20031.
- 1250 KRAPF, D., VISCONTI, P. E., ARRANZ, S. E. & CABADA, M. O. (2007). Egg water from the
1251 amphibian *Bufo arenarum* induces capacitation-like changes in homologous spermatozoa.
1252 *Developmental Biology* **306**, 516–524.
- 1253 KRAPF, D., O'BRIEN, E. D., CABADA, M. O., VISCONTI, P. E. & ARRANZ, S. E. (2009). Egg water

- 1254 from the amphibian *Bufo arenarum* modulates the ability of homologous sperm to undergo
 1255 the acrosome reaction in the presence of the vitelline envelope. *Biology of Reproduction* **80**,
 1256 311–319.
- 1257 KRAPF, D., O'BRIEN, E., MAIDAGÁN, P. M., MORALES, E. S., VISCONTI, P. E. & ARRANZ, S. E.
 1258 (2014). Calcineurin regulates progressive motility activation of *Rhinella (Bufo) arenarum*
 1259 sperm through dephosphorylation of PKC substrates. *Journal of Cellular Physiology* **229**,
 1260 1378–1386.
- 1261 KUBO-IRIE, M., IRIE, M., NAKAZAWA, T. & MOHRI, H. (2003). Ultrastructure and function of long
 1262 and short sperm in Cicadidae (Hemiptera). *Journal of Insect Physiology* **49**, 983–991.
- 1263 KUZAN, F. B., FLEMING, A. D. & SEIDEL, G. E. JR. (1984). Successful fertilization *in vitro* of fresh
 1264 intact oocytes by perivitelline (acrosome reacted) spermatozoa of the rabbit. *Fertility*
 1265 *and Sterility* **41**, 766–770.
- 1266 LAFLAMME, B. A., RAVI RAM, K. & WOLFNER, M. F. (2012). The *Drosophila melanogaster*
 1267 seminal fluid protease “seminase” regulates proteolytic and post-mating reproductive
 1268 processes. *PLoS Genetics* **8**, e1002435.
- 1269 LAHNSTEINER, F., WEISMANN, T. & PATZNER, R. A. (1995). Composition of the ovarian fluid in 4
 1270 salmonid species - *Oncorhynchus mykiss*, *Salmo trutta lacusris*, *Salvelinus alpinus* and
 1271 *Hucho hucho*. *Reproduction Nutrition Development* **35**, 465–474.
- 1272 LAMBERT, C. C. (1982). The ascidian sperm reaction. *American Zoologist* **22**, 841–849.
- 1273 LA SPINA, F., PUGA MOLINA, L. C., ROMAROWSKI, A., VITALE, A. M., FALZONE, T. L., KRAPF,
 1274 D., HIROHASHI, N. & BUFFONE, M. G. (2016). Mouse sperm begin to undergo acrosomal
 1275 exocytosis in the upper isthmus of the oviduct. *Developmental Biology* **411**, 172–182.
- 1276 LIBERTI, J., BAER, B. & BOOMSMA, J. J. (2016). Queen reproductive tract secretions enhance

- 1277 sperm motility in ants. *Biology Letters* **12**, 20160722.
- 1278 LIBERTI, J., BAER, B. & BOOMSMA, J. J. (2018). Rival seminal fluid induces enhanced sperm
1279 motility in a polyandrous ant. *BMC Evolutionary Biology* **18**, 28.
- 1280 LIN, M. & RODGER, J.C. (1999). Acrosome formation during sperm transit through the
1281 epididymis in two marsupials, the tammar wallaby (*Macropus eugenii*) and the brushtail
1282 possum (*Trichosurus vulpecula*). *Journal of Anatomy* **194**, 223–232.
- 1283 LIU, H. & KUBLI, E. (2003). Sex-peptide is the molecular basis of the sperm effect in *Drosophila*
1284 *melanogaster*. *Proceedings of the National Academy of Sciences, USA* **100**, 9929–9933.
- 1285 LONGO, G., SOTTILE, L., VISCUSO, R., GIUFFRIDA, A. & PRIVITERA, R. (1993). Ultrastructural
1286 changes in sperm of *Eyprepocnemis plorans* (Charpentier) (Orthoptera: Acrididae) during
1287 storage of gametes in female genital tract. *Invertebrate Reproduction and Development* **24**,
1288 1–6.
- 1289 LUPETTI, P., MERCATI, D. & DALLAI, R. (2001). The sperm glycocalyx of *Pezotettix giornai*
1290 (Rossi) (Insecta: Orthoptera) after quick-freeze, deep-etching. *Italian Journal of Anatomy*
1291 *and Embryology* **106 (Suppl 2)**, 181–188.
- 1292 LÜPOLD, S., MANIER, M. K., PUNIAMOORTHY, N., SCHOFF, C., STARMER, W. T., LUEPOLD, S. H.,
1293 BELOTE, J. M. & PITNICK, S. (2016). How sexual selection can drive the evolution of costly
1294 sperm ornamentation. *Nature* **533**, 535–538.
- 1295 MAGAREY, G. M. & MATE, K. E. (2003). Fertilization following intracytoplasmic sperm injection
1296 of *in vivo* and *in vitro* matured oocytes from an Australian marsupial, the tammar wallaby
1297 (*Macropus eugenii*). *Zygote* **11**, 339–346.
- 1298 MAKIELSKI, S. K. (1966). The structure and maturation of the spermatozoa of *Sciara coprophila*.
1299 *Journal of Morphology* **118**, 11–42.

- 1300 MANN, T. (1984). *Spermatophores*. Springer Verlag: New York.
- 1301 MANN, T., MARTIN, A. W. JR. & THIERSCH, J. B. (1970). Male reproductive tract, spermatophores
1302 and spermatophoric reaction in the giant octopus of the North Pacific, *Octopus dofleini*
1303 *martini*. *Proceedings of the Royal Society, Series B* **175**, 31–61.
- 1304 MANNING A. (1967). The control of sexual receptivity in female *Drosophila*. *Animal Behaviour*
1305 **15**, 239–250.
- 1306 MATE, K. E. & RODGER, J. C. (1996). Capacitation and the acrosome reaction in marsupial
1307 spermatozoa. *Reproduction Fertility and Development* **8**, 595–603.
- 1308 MATE, K. E., SIDHU, K. S., MOLINIA, F. C., GLAZIER, A. M. & RODGER, J. C. (2000). Sperm
1309 binding and penetration of the zona pellucida *in vitro* but not sperm-egg fusion in an
1310 Australian marsupial, the brushtail possum (*Trichosurus vulpecula*). *Zygote* **8**, 189–196.
- 1311 MATZKE-KARASZ, R., SMITH, R. J. & HEB, M. (2017). Removal of extracellular coat from giant
1312 sperm in female receptacle induces motility in *Mytilocyris mytiloides* (Cyprididae, Ostracoda,
1313 Crustacea). *Cell and Tissue Research* **368**, 171–186.
- 1314 MICHALIK, P. (2007). Spermatozoa and spermiogenesis of *Liphistius* cf. *phuketensis*
1315 (Mesothelae, Araneae, Arachnida) with notes on phylogenetic implications. *Arthropod*
1316 *Structure and Development* **36**, 327–335.
- 1317 MICHALIK, P., HAUPT, J. & ALBERTI, G. (2004). On the occurrence of coenospermia in
1318 mesothelid spiders (Araneae: Heptathelidae). *Arthropod Structure and Development* **33**, 173–
1319 181.
- 1320 MICHALIK, P., REIHER, W., TINTELNOT-SUHM, M., COYLE, F. A. & ALBERTI, G. (2005). Female
1321 genital system of the folding-trapdoor spider *Antrodiaetus unicolor* (Hentz, 1842)

- 1322 (Antrodiaetidae, Araneae): Ultrastructural study of form and function with notes on
 1323 reproductive biology of spiders. *Journal of Morphology* **263**, 284–309.
- 1324 MIRANDA, P. V., ALLAIRE, A., SOSNIK, J. & VISCONTI, P. E. (2009). Localization of low-density
 1325 detergent-resistant membrane proteins in intact and acrosome-reacted mouse sperm. *Biology*
 1326 *of Reproduction* **80**, 897–904.
- 1327 MONCLUS, M. A. & FORNES, M. W. (2016). Sperm conjugation in mammal reproductive
 1328 function: different names for the same phenomenon? *Molecular Reproduction and*
 1329 *Development* **83**, 884–896.
- 1330 MOORE, H. D. & TAGGART, D. A. (1993). *In vitro* fertilization and embryo culture in the grey
 1331 short-tailed opossum, *Monodelphis domestica*. *Journal of Reproduction and Fertility* **98**,
 1332 267–274.
- 1333 MORROW, E. H. (2004). How the sperm lost its tail: the evolution of aflagellate sperm. *Biological*
 1334 *Reviews* **79**, 795–814.
- 1335 MOTHES, U. & SEITZ, K.-A. (1981). The transformation of male sex cells of *Tetranychus urticae*
 1336 K (Acari, Tetranychidae) during passage from the testis to the oocytes: an electron
 1337 microscopic study. *International Journal of Invertebrate Reproduction* **4**, 81–94.
- 1338 MURO, Y., HASUWA, H., ISOTANI, A., MIYATA, H., YAMAGATA, K., IKAWA, M., YANAGIMACHI, R.
 1339 & OKABE, M. (2016). Behavior of mouse spermatozoa in the female reproductive tract from
 1340 soon after mating to the beginning of fertilization. *Biology of Reproduction* **94**, 80.
- 1341 NAKANISHI, T., IKAWA, M., YAMADA, S., PARVINEN, M., BABA, T., NISHIMUNE, Y. & OKABE, M.
 1342 (1999). Real-time observation of acrosomal dispersal from mouse sperm using GFP as a
 1343 marker protein. *FEBS Letters* **449**, 277–283.

- 1344 NDIAYE, M., MATTEI, X. & THIAW, O. T. (1997). Maturation of mosquito spermatozoa during
 1345 their transit throughout the male and female reproductive systems. *Tissue & Cell* **6**, 675–678.
- 1346 NEWPORT, G. (1851). On the impregnation of the ovum in the amphibian. *Philosophical*
 1347 *Transactions of the Royal Society of London (First Series Part 1)* **141**, 169–242.
- 1348 NIXON, B., ANDERSON, A. L., SMITH, N. D., MCLEOD, R. & JOHNSTON, S. D. (2016a). The
 1349 Australian saltwater crocodile (*Crocodylus porosus*) provides evidence that the capacitation
 1350 of spermatozoa may extend beyond the mammalian lineage. *Proceedings of the Royal*
 1351 *Society, Series B* **283**, 20160495.
- 1352 NIXON, B., ECROYD, H. W., DACHEUX, J.-L. & JONES, R. C. (2011). Monotremes provide a key to
 1353 understanding the evolutionary significance of epididymal sperm maturation. *Journal of*
 1354 *Andrology* **32**, 665–671.
- 1355 NIXON, B., ECROYD, H. W., DACHEUX, J.-L., DACHEUX, F., LABAS, V., JOHNSTON, S. D. & JONES,
 1356 R. C. (2016b). Formation and dissociation of sperm bundles in monotremes. *Biology of*
 1357 *Reproduction* **95**, **91**, 1–11.
- 1358 NIXON, B., EWEN, K. A., KRIVANEK, K. M., CLULOW, J., KIDD, G., ECROYD, H. W. & JONES, R. C.
 1359 (2014). Post-testicular sperm maturation and identification of an epididymal protein in the
 1360 Japanese quail (*Coturnix coturnix japonica*). *Reproduction* **147**, 265–277.
- 1361 NIXON, B., JOHNSTON, S. D., SKERRETT-BYRNE, D. A., ANDERSON, A. L., STANGER, S. J.,
 1362 BROMFIELD, E. G., MARTIN, J. H., HANSBRO, P. M. & DUN, M. D. (2019b). Modification of
 1363 crocodile spermatozoa refutes the tenet that post-testicular sperm maturation is restricted to
 1364 mammals. *Molecular and Cellular Proteomics* **18**, S59-S76.
- 1365 Ó FOIGHIL, D. (1985a). Fine structure of *Lasaea subviridis* and *Mysella tumida* sperm (Bivalvia:
 1366 Galeommatacea). *Zoomorphology* **105**, 125–132.

- 1367 Ó FOIGHIL, D. (1985*b*). Sperm transfer and storage in the brooding bivalve *Mysella tumida*.
 1368 *Biological Bulletin* **169**, 602–614.
- 1369 OHTO, U., ISHIDA, H., KRAYUKHINA, E., UCHIYAMA, S., INOUE, N. & SHIMIZU, T. (2016).
 1370 Structure of IZUMO1-JUNO reveals sperm-oocyte recognition during mammalian
 1371 fertilization. *Nature* **534**, 566–569.
- 1372 OKABE, M. (2014). Mechanism of fertilization: a modern view. *Experimental Animals* **63**, 357–
 1373 365.
- 1374 OKABE, M., ADACHI, T., TAKADA, K., ODA, H., YAGASAKI, M., KOHAMA, Y. & MIMURA, T.
 1375 (1987). Capacitation-related changes in antigen distribution on mouse sperm heads and its
 1376 relation to fertilization rate *in vitro*. *Journal of Reproductive Immunology* **11**, 91–100.
- 1377 OLIVER, J. H. JR. & BRINTON, L. P. (1971). Sperm maturation in ticks: an example of capacitation
 1378 in invertebrates? *Proceedings of the Third International Congress of Acarology*. W. Junk,
 1379 The Hague, pp. 733–737.
- 1380 O’RAND, M. G. (1972). *In vitro* fertilization and capacitation-like interaction in the hydroid
 1381 *Campanularia flexuosa*. *Journal of Experimental Zoology* **182**, 299–305.
- 1382 O’RAND, M. G. (1974). Gamete interaction during fertilization in *Campanularia* - The female
 1383 epithelial cell surface. *American Zoologist* **14**, 487–493.
- 1384 O’RAND, M. G. & MILLER, R. L. (1974). Spermatozoan vesicle loss during penetration of the
 1385 female gonangium in the hydroid *Campanularia flexuosa*. *Journal of Experimental Zoology*
 1386 **188**, 179–193.
- 1387 OSANAI, M. & ISONO, M. (1997). Dissociation of eusperm bundles by acids, especially by
 1388 succinate accumulated in the spermatophore of the silkworm, *Bombyx mori*. *Invertebrate*
 1389 *Reproduction and Development* **31**, 99–108.

- 1390 OSANAI, M., KASUGA, H. & AIGAKI, T. (1989a). Induction of motility of apyrene sperm bundles
 1391 of the silkmoth, *Bombyx mori*, by initiatorin and trypsin. *Invertebrate Reproduction and*
 1392 *Development* **15**, 97–103.
- 1393 OSANAI, M., KASUGA, H. & AIGAKI, T. (1989b). Isolations of eupyrene sperm bundles and
 1394 apyrene spermatozoa from seminal fluid of the silkmoth, *Bombyx mori*. *Journal of Insect*
 1395 *Physiology* **35**, 401–408.
- 1396 PEMBERTON, A. J., NOBLE, L. R. & BISHOP, J. D. D. (2003). Frequency dependence in matings
 1397 with water-borne sperm. *Journal of Evolutionary Biology* **16**, 289–301.
- 1398 PENG, J., CHEN, S., BÜSSER, S., LIU, H., HONEGGER, T. & KUBLI, E. (2005). Gradual release of
 1399 sperm bound sex-peptide controls female postmating behavior in *Drosophila*. *Current*
 1400 *Biology* **15**, 207–213.
- 1401 PHILLIPS, D. M. (1966a). Fine structure of *Sciara coprophila* sperm. *Journal of Cell Biology* **30**,
 1402 499–517.
- 1403 PHILLIPS, D. M. (1966b). Observations on spermiogenesis in the fungus gnat *Sciara coprophila*
 1404 sperm. *Journal of Cell Biology* **30**, 477–497.
- 1405 PICARD, A. (1980). Spermatogenesis and sperm-spermatheca relations in *Spirorbis spirorbis* (L.).
 1406 *International Journal of Invertebrate Reproduction* **2**, 73–83.
- 1407 PINCUS, G. & ENZMANN, E. V. (1934). Can mammalian eggs undergo normal development in
 1408 vitro? *Proceedings of the National Academy of Sciences, USA* **20**, 121–122.
- 1409 PITNICK, S., MARKOW, T. A. & SPICER, G. S. (1995a). Delayed male maturity is a cost of
 1410 producing large sperm in *Drosophila*. *Proceedings of the National Academy of Sciences,*
 1411 *USA* **92**, 10614–10618.

- 1412 PITNICK, S., MARKOW, T. A. & SPICER, G. S. (1999). Evolution of multiple kinds of female
1413 sperm-storage organs in *Drosophila*. *Evolution* **53**, 1804–1822.
- 1414 PITNICK, S., SPICER, G. S. & MARKOW, T. A. (1995*b*). How long is a giant sperm? *Nature* **375**,
1415 109.
- 1416 PITNICK, S., WOLFNER, M. F. & SUAREZ, S.S. (2009*b*). Ejaculate-female and sperm-female
1417 interactions. In: *Sperm Biology: An Evolutionary Perspective* (TR Birkhead, DJ Hosken and
1418 S Pitnick, eds.), pp. 247–304. Academic Press, London.
- 1419 POLAND, V., EUBEL, H., KING, M., SOLHEIM, C., MILLAR, A. H. & BAER, B. (2011). Stored sperm
1420 differs from ejaculated sperm by proteome alterations associated with energy metabolism in
1421 the honeybee *Apis mellifera*. *Molecular Ecology* **20**, 2643–2654.
- 1422 RAVI RAM, K. & WOLFNER, M.F. (2009). A network of interactions among seminal proteins
1423 underlies the long-term postmating response in *Drosophila*. *Proceedings of the National*
1424 *Academy of Sciences, USA* **106**, 15384–15389.
- 1425 REGER, J. F. (1961). The fine-structure of spermatids from the tick, *Amblyomma dissimili*.
1426 *Journal of Ultrastructure Research* **5**, 584–599.
- 1427 REGER, J. F. (1962). A fine-structure study on spermiogenesis in the tick, *Amblyomma dissimili*
1428 with special reference to the development of motile processes. *Journal of Ultrastructure*
1429 *Research* **7**, 550–565.
- 1430 REGER, J. F. (1963). Spermiogenesis in the tick, *Amblyomma dissimili*, as revealed by electron
1431 microscopy. *Journal of Ultrastructure Research* **8**, 607–621.
- 1432 REGER, J. F. (1970). Some aspects of the fine structure of filiform spermatozoa (ostracod,
1433 *Cypridopsis* sp.) lacking tubule sub-structure. In: *Comparative Spermatology* (Ed. by B.
1434 Baccetti), pp. 237–245. New York: Academic Press.

- 1435 REGER, J. F. (1974). The origin and fine structure of cellular processes in spermatozoa of the tick,
 1436 *Dermacentor andersoni*. *Journal of Ultrastructure Research* **48**, 420–434.
- 1437 RENIERI, T. & TALLURI, M. V. (1974). Sperm modification in the female ducts of a grasshopper.
 1438 *Monitore Zoologico Italiano* **8**, 1–9.
- 1439 RICHINGS, N. M., SHAW, G., TEMPLE-SMITH, P. D. & RENFREE, M. B. (2004). Intra-cytoplasmic
 1440 sperm injection in a marsupial. *Reproduction* **128**, 595–605.
- 1441 RIEMANN, J. G. (1970). Metamorphosis of sperm of the cabbage looper, *Trichoplusia ni*, during
 1442 passage from the testes to the female spermathecae. pp. 321–331, In: *Comparative*
 1443 *Spermatology* (Baccetti B, ed), Accademia Nazionale Dei Lincei, Quaderono N. 137.
- 1444 RIEMANN, J. G. & GIEBULTOWICZ, J. M. (1991). Secretion in the upper vas deferens of the gypsy
 1445 moth correlated with the circadian rhythm of sperm release from the testes. *Journal of Insect*
 1446 *Physiology* **37**, 53–62.
- 1447 RIEMANN, J. G. & GIEBULTOWICZ, J. M. (1992). Sperm maturation in the vasa deferentia of the
 1448 gypsy-moth, *Lymantria dispar* (Lepidoptera, Lymantriidae). *Internaltional Journal of Insect*
 1449 *Morphology andf Embryology* **21**, 271–284.
- 1450 RIEMANN, J. G. & THORSON B. J. (1971). Sperm maturation in the male and female genital tracts
 1451 of *Anagasta kühniella* (Lepidoptera: Pyralididae). *Internaltional Journal of Insect*
 1452 *Morphology andf Embryology* **1**, 11–19.
- 1453 ROBISON, W. G. (1970). Unusual arrangement of microtubules in relation to mechanisms of
 1454 sperm movement. In: *Comparative Spermatology* (Ed. by B. Baccetti), pp. 311–320. New
 1455 York: Academic Press.
- 1456 RODGER, J. C. (1994). Prefertilization gamete maturation events in marsupials. *Reproduction*
 1457 *Fertility and Development* **6**, 473–483.

- 1458 RODGER, J. C. & BEDFORD, J. M. (1982). Separation of sperm pairs and sperm-egg interaction in
 1459 the opossum, *Virginia didephis*. *Journal of Reproduction and Fertility* **64**, 171–179.
- 1460 ROSENGRAVE, P., TAYLOR, H., MONTGOMERIE, R., METCALF, V., MCBRIDE, K. & GEMMELL, N. J.
 1461 (2009). Chemical composition of seminal and ovarian fluids of chinook salmon
 1462 (*Oncorhynchus tshawytscha*) and their effects on sperm motility traits. *Comparative*
 1463 *Biochemistry and Physiology, Part A* **152**, 123–129.
- 1464 ROTHSCILD, L. (1961). Structure and movements of tick spermatozoa (Arachnida: Acari).
 1465 *Quarterly Journal of Microscopic Science* **102**, 239–247.
- 1466 SAHARA, K. & TAKEMURA, Y. (2003). Application of artificial insemination technique to
 1467 eupyrene and/or apyrene sperm in *Bombyx mori*. *Journal of Experimental Zoology* **297A**,
 1468 196–200.
- 1469 SALING, P. M. & STOREY, B. T. (1979). Mouse gamete interaction during fertilization *in vitro*:
 1470 chlortetracycline as a fluorescent probe for the mouse acrosome reaction. *Journal of Cell*
 1471 *Biology* **83**, 544–555.
- 1472 SASAKAWA, K. (2007). Sperm bundle and reproductive organs of carabid beetles tribe
 1473 Pterostichini (Coleoptera: Carabidae). *Naturwissenschaften* **94**, 384–391.
- 1474 SASTRY, A.N. (1979). Pelecypoda (excluding Ostreidae). Pp. 113–292 in *Reproduction of*
 1475 *Marine Invertebrates, Vol. V*. AC Giese and JS Pearse, eds. New York: Academic Press.
- 1476 SCHUBERT, L. F., KRÜGER, S., MORITZ, G. B. & SCHUBERT, V. (2017). Male reproductive system
 1477 and spermatogenesis of *Limodromus assimilis* (Paykull 1790). *PLoS One* **12**, e0180492.
- 1478 SELMI, M. G., BIGLIARDI, E. & GIUSTI, F. (1989). Morphological modifications in stored
 1479 heterospermatozoa of *Oxyloma elegans* (Pulmonata: Stylommatophora). *Journal of*
 1480 *Ultrastructure and Molecular Structure Research* **102**, 82–86.

- 1481 SETIADI, D., LIN, M. & RODGER, J. C. (1997). Posttesticular development of spermatozoa of the
1482 tammar wallaby (*Macropus eugenii*). *Journal of Anatomy* **190**, 275–288.
- 1483 SEVER, D. M. (2002). Female sperm storage in amphibians. *Journal of Experimental Zoology*
1484 **292**, 165–179.
- 1485 SHEPHERD, J., LEVINE, S. & HALL, J. D. (1982a). Maturation of tick spermatozoa *in vitro*.
1486 *International Journal of Invertebrate Reproduction* **4**, 311–321.
- 1487 SHEPHERD, J., OLIVER, J. H. JR. & HALL, J. D. (1982b). A polypeptide from male accessory
1488 glands which triggers maturation of tick spermatozoa. *International Journal of Invertebrate*
1489 *Reproduction* **5**, 129–137.
- 1490 SHIVERS, C. A. & JAMES, J. M. (1970a). Capacitation of frog sperm. *Nature* **227**, 183–184.
- 1491 SHIVERS, C. A. & JAMES, J. M. (1970b). Morphology and histochemistry of the oviduct and egg-
1492 jelly layers in the frog, *Rana pipiens*. *Anatomical Record* **166**, 541–556.
- 1493 SHIVERS, C. A. & JAMES, J. M. (1971). Fertilization of antiserum-inhibited frog eggs with
1494 “capacitated” sperm. *Biology of Reproduction* **5**, 229–235.
- 1495 SIDHU, K. S., MATE, K. E., MOLINIA, F. C., GLAZIER, A. M. & RODGER, J. C. (1999a). Secretory
1496 proteins from the female reproductive tract of the brushtail possum (*Trichosurus vulpecula*):
1497 binding to sperm and effects on sperm survival *in vitro*. *Reproduction Fertility and*
1498 *Development* **11**, 329–336.
- 1499 SIDHU, K. S., MATE, K. E., MOLINIA, F. C. & RODGER, J. C. (1999b). Induction of thumbtack
1500 sperm during coculture with oviduct epithelial cell monolayers in a marsupial, the brushtail
1501 possum (*Trichosurus vulpecula*). *Biology of Reproduction* **61**, 1356–1361.
- 1502 SILBERGLIED, D. R., SHEPHERD, J. G. & DICKINSON, J. L. (1984). Eunuchs: the role of apyrene
1503 sperm in Lepidoptera? *The American Naturalist* **12**, 255–265.

- 1504 SIMMONS, L. W., ROBERTS, J. D. & DZIMINSKI, M. A. (2009). Egg jelly influences sperm motility
1505 in the externally fertilizing frog, *Crinia georgiana*. *Journal of Evolutionary Biology* **22**, 225–
1506 229.
- 1507 SINGH, A., BUEHNER, N. A., LIN, H., BARANOWSKI, K. J., FINDLAY, G. D. & WOLFNER, M. F.
1508 (2018). Long-term interaction between *Drosophila* sperm and sex peptide is mediated by
1509 other seminal proteins that bind only transiently to sperm. *Insect Biochemistry and Molecular*
1510 *Biology* **102**, 43-51.
- 1511 SISTINA, Y., LIN, M., MATE, K. E., ROBINSON, E. S. & RODGER, J. C. (1993a). The unique stability
1512 of the marsupial sperm acrosomal membranes examined by unprotected freeze-thawing and
1513 treatment with the detergent Triton X-100. *Reproduction Fertility and Development* **5**, 1–14.
- 1514 SISTINA, Y., LIN, M., MATE, K. E. & RODGER, J. C. (1993b). Induction of the marsupial acrosome
1515 reaction *in vitro* by treatment with diacylglycerols. *Journal of Reproduction and Fertility* **99**,
1516 335–341.
- 1517 SMITH, R. J., MATZKE-KARASZ, R., KAMIYA, T. & DE DECKKER, P. (2016). Sperm lengths of non-
1518 marine cypridoidean ostracods (Crustacea). *Acta Zoologica* **97**, 1–17.
- 1519 SNOOK, R. R. & KARR, T. L. (1998). Only long sperm are fertilization-competent in six sperm-
1520 heteromorphic *Drosophila* species. *Current Biology* **8**, 291–294.
- 1521 STOCKLEY, P., GAGE, M. J. G., PARKER, G. A. & MØLLER, A. P. (1997). Sperm competition in
1522 fishes: the evolution of testis size and ejaculate characteristics. *The American Naturalist* **149**,
1523 933–954.
- 1524 SUAREZ, S. S. (2002). Formation of a reservoir of sperm in the oviduct. *Reproduction in*
1525 *Domestic Animals* **37**, 140–143.
- 1526 SUAREZ, S. S. (2008). Control of hyperactivation in sperm. *Human Reproduction Update* **14**,

- 1527 647–657.
- 1528 SUAREZ, S. S. (2016). Mammalian sperm interactions with the female reproductive tract. *Cell*
- 1529 *and Tissue Research* **363**, 185–194
- 1530 SUAREZ, S. S. & PACEY A. A. (2006). Sperm transport in the female reproductive tract.
- 1531 *Human Reproduction Update* **12**, 23–37.
- 1532 SWALLOW, J. G. & WILKINSON, G. S. (2002). The long and short of sperm polymorphisms in
- 1533 insects. *Biological Reviews* **77**, 153–182.
- 1534 TAKAMI, Y. & SOTA, T. (2007). Sperm competition promotes diversity of sperm bundles in
- 1535 *Ohomopterus* ground beetles. *Naturwissenschaften* **94**, 543–550.
- 1536 TAKEMURA, Y., SAHARA, K., MOCHIDA, Y. & OHNUMA, A. (2006). Apyrene sperm from the
- 1537 triploid donors restore fecundity of cryopreserved semen in *Bombyx mori*. *Journal of Insect*
- 1538 *Physiology* **52**, 1021–1026.
- 1539 TEMKIN, M. H. (1994). Gamete spawning and fertilization in the gymnolaemate bryozoan
- 1540 *Membranipora membranacea*. *Biological Bulletin* **187**, 143–155.
- 1541 TEMKIN, M. H. & BORTOLAMI, S. B. (2004). Waveform dynamics of spermatzeugmata during
- 1542 the transfer from paternal to maternal individuals of *Membranipora membranacea*.
- 1543 *Biological Bulletin* **206**, 35–45.
- 1544 THALER, C. D., MIYATA, H., HAIMO, L. T. & CARDULLO, R. A. (2013). Waveform generation is
- 1545 controlled by phosphorylation and swimming direction is controlled by Ca^{2p} in sperm from
- 1546 the mosquito *Culex quinquefasciatus*. *Biology of Reproduction* **89**, 135.
- 1547 TILL-BOTTRAUD, I., JOLY, D., LACHAISE, D. & SNOOK, R. R. (2005). Pollen and sperm
- 1548 heteromorphism: convergence across kingdoms? *Journal of Evolutionary Biology* **18**, 1–18.

- 1549 TOSTI, E., DI COSMO, A., CUOMO, A., DI CRISTO, C. & GRAGNANIELLO, G. (2001). Progesterone
 1550 induces activation in *Octopus vulgaris* spermatozoa. *Molecular Reproduction and*
 1551 *Development* **59**, 97–105.
- 1552 TURNER, E. & MONTGOMERIE, R. (2002). Ovarian fluid enhances sperm movement in Arctic
 1553 charr. *Journal of Fish Biology* **60**, 1570–1579.
- 1554 UHL, G. (1994). Ultrastructure of the accessory glands in female genitalia of *Pholcus*
 1555 *phalangioides* (Fuesslin, 1771) (Pholcidae; Araneae). *Acta Zoologica Stockholm* **75**, 13–25.
- 1556 UHL, G. (1996). Sperm storage secretion of female cellar spiders (*Pholcus phalangioides*;
 1557 Araneae): a gel-electrophoretic analysis. *Journal of Zoology, London* **240**, 153–161.
- 1558 UHL, G. (2000). Two distinctly different sperm storage organs in female *Dysdera erythrina*
 1559 (Araneae: Dysderidae). *Arthropod Structure and Development* **29**, 163–169.
- 1560 URSPRUNG, H. & SCHABTACH, E. (1965). Fertilization in tunicates: Loss of the paternal
 1561 mitochondrion prior to sperm entry. *Journal of Experimental Zoology* **159**, 379–384.
- 1562 VISCUSO, R., NARCISI, L., SOTTILE, L. & BARONE, N. (1998). Structure of spermatodesms of
 1563 Orthoptera Tettinonioidae. *Tissue & Cell* **30**, 453–463.
- 1564 VISCUSO, R., NARCISI, L., SOTTILE, L. & VIOLETTA BRUNDO, M. (2001). Role of male accessory
 1565 glands in spermatodesm reorganization in Orthoptera Tettigonioidae. *Tissue & Cell* **33**, 33–
 1566 39.
- 1567 VISCUSO, R., VIOLETTA BRUNDO, M. & SOTTILE, L. (2002). Mode of transfer of spermatozoa in
 1568 Orthoptera Tettigonioidae. *Tissue & Cell* **34**, 337–348.
- 1569 VÖCKING, O., UHL, G. & MICHALIK, P. (2013). Sperm dynamics in spiders (Araneae):
 1570 ultrastructural analysis of the sperm activation process in the garden spider *Argiope*
 1571 *bruennichi* (Scopoli, 1772). *PLoS ONE* **8(9)**, e72660.

- 1572 WAKE, M. H. & DICKIE, R. (1998). Oviductal structure and function and reproductive modes in
1573 amphibians. *Journal of Experimental Zoology* **282**, 477–506.
- 1574 WATANABE, T., ITOH, T., WATANABE, A. & ONITAKE, K. (2003). Characteristics of sperm
1575 motility induced on the egg-jelly in the internal fertilization of the newt, *Cynops*
1576 *pyrrhogaster*. *Zoological Science* **20**, 345–342.
- 1577 WATANABE, T., KUBO, H., TAKESHIMA, S., NAKAGAWA, M., OHTA, M., KAMIMURA, S.,
1578 TAKAYAMA-WATANABE, E., WATANABE, A. & ONITAKE, K. (2010). Identification of the
1579 sperm motility-initiating substance in the newt, *Cynops pyrrhogaster*, and its possible
1580 relationship with the acrosome reaction during internal fertilization. *International Journal of*
1581 *Developmental Biology* **54**, 591–597.
- 1582 WATANABE, A. & ONITAKE, K. (2002). The urodele egg-coat as the apparatus adapted for the
1583 internal fertilization. *Zoological Science* **19**, 1341–1347.
- 1584 WATNICK, T. J., JIN, Y., MATUNIS, E., KERNAN, M. J. & MONTELL, C. (2003). A flagellar
1585 polycystin-2 homolog required for male fertility in *Drosophila*. *Current Biology* **13**, 2179–
1586 2184.
- 1587 WEBBER, H. H. (1977). Gastropoda: Prosobranchia. Pp. 1–97 in *Reproduction of Marine*
1588 *Invertebrates, Vol. IV*. AC Giese and JS Pearse, eds. New York: Academic Press.
- 1589 WESTHEIDE, W. (1988). The ultrastructure of the spermatozoon in *Pisioine remota* (Annelida:
1590 Polychaeta) and its transformation in the receptaculum seminis. *Journal of Submicroscopic*
1591 *Cytology and Pathology* **20**, 169–178.
- 1592 WILLIAMS, M., BARRATT, C. L., HILL, C. J., WARREN, M. A., DUNPHY, B. & COOKE, I. D. (1992).
1593 Recovery of artificially inseminated spermatozoa from the fallopian tubes of a woman
1594 undergoing total abdominal hysterectomy. *Human Reproduction* **7**, 506–509

- 1595 WINGSTRAND, K. G. (1988). Comparative spermatology of the Crustacea Entomostraca. 2.
 1596 Subclass Ostracoda. *Biologiske Skrifter Kongelige Danske Videnskabernes Selskab* **32**, 1–
 1597 149.
- 1598 YANAGIMACHI, R. (1982). Requirement of extracellular calcium ions for various stages of
 1599 fertilization and fertilization-related phenomena in the hamster. *Gamete Research* **5**, 323–
 1600 344.
- 1601 YANG, Y. & LU, X. (2011). *Drosophila* sperm motility in the reproductive tract. *Biology of*
 1602 *Reproduction* **84**, 1005–1015.
- 1603 YAPICI, N., KIM, Y. J., RIBEIRO, C. & DICKSON, B. J. (2008). A receptor that mediates the post-
 1604 mating switch in *Drosophila* reproductive behaviour. *Nature* **451**, 33–37.
- 1605 YASUZUMI, G. (1979). Spermatogenesis in animals as revealed by electron microscopy. Some
 1606 modifications of cell surface in developing spermatids of the grasshopper. *Monitore*
 1607 *Zoologico Italiano* **13**, 265–277.
- 1608 ZHANG, L., HAN, X. K., QI, Y. Y., LIU, Y. & CHEN Q. S. (2008). Seasonal effects on apoptosis and
 1609 proliferation of germ cells in the testes of the Chinese soft-shelled turtle, *Pelodiscus sinensis*.
 1610 *Theriogenology* **69**, 1148–1158.
- 1611 ZHANG, L., YANG, P., BIAN, X., ZHANG, Q., ULLAH, S., WAQAS, Y., CHEN, X., LIU, Y., CHEN, W.,
 1612 LE, Y., CHEN, B., WANG, S. & CHEN, Q. (2015). Modification of sperm morphology during
 1613 long-term sperm storage in the reproductive tract of the Chinese soft-shelled turtle,
 1614 *Pelodiscus sinensis*. *Scientific Reports* **5**, 16096.
- 1615